

**LECTURE NOTES**  
**On**  
**ENGINEERING MATHEMATICS-II**  
**2<sup>nd</sup> Semester**

**Diploma in Electrical and Electronics Engineering**



**Prepared By:**  
**Sushreesangita Rout**

**Department of Electrical and Electronics Engineering**  
**Vedang Institute of Technology**  
**Bhubaneswar, Khurda**

## TABLE OF CONTENTS

Chapter no	Topics	Sub Topics	Page no
1	Vector Algebra	Introduction Representation of vector Magnitude and direction of vector Types of vectors Operations on vectors (addition, subtraction and scalar multiplication) Position Vector Representation of vector in component form (2-D and 3-D) Scalar or dot product of two vectors Geometrical meaning of dot product Scalar and Vector projection of two vectors Angle between two vectors Vector product Geometrical meaning of vector product (Area of triangle and parallelogram)	1-18
2	Limits and Continuity	Introduction Definition of function, based on set theory Domain and Range of a function Classification of function Types of functions (constant, Identity, $a^x, \log x$ [x], trigonometric, etc) Introduction to limit Existence of limit Method of evaluation of limit Evaluation of infinite limits Evaluation of limits using formulas and substitution method Continuity of a function at a point	19-40

3	Derivatives	<p>Introduction  Derivative of a function at a point  Geometrical Interpretation of derivative  Derivative of some standard functions <math>x^n, a^x</math>  etc.  Derivative of composite functions (chain rule)  Method of differentiation using substitution  methods and logarithms  Differentiation of parametric functions  Differentiation of a function w.r.t. another  function.  Differentiation of Implicit functions  Application of derivatives  Successive Differentiation  Partial Differentiation (function of  two variables up to second order)</p>	41-77
4	Integration	<p>Definition of integration as inverse of  differentiation  Integrals of some standard function  Methods of  Integration i) <math>\int \frac{dx}{x^2+a^2}, \int \frac{dx}{x^2+a^2}</math>  Substitution  Integration of forms etc using substitution method  ii) Integration By parts  <math>\int \sqrt{a^2 - x^2} dx, \int \sqrt{a^2 + x^2} dx</math> Integration  of forms using by parts  Definition of Definite Integral  Fundamental theorem of Integral Calculus  Properties of Definite Integral  Application of Integration  i) Area enclosed by a curve and X- axis. ii)  Area of a circle with centre at origin.</p>	61-77
5	Differential Equation	<p>Order and degree of a differential equation  Solution of Differential equation 1<sup>st</sup> order and  1<sup>st</sup> degree  i) variable separable  method <math>\frac{dy}{dx} + Py = Q,</math>  ii) Linear equation where P,Q are functions of  x.</p>	78-86

# UNIT-I

## VECTORS

### INTRODUCTION:-

In our real life situation we deal with physical quantities such as distance, speed, temperature, volume etc. These quantities are sufficient to describe change of position, rate of change of position, body temperature or temperature of a certain place and space occupied in a confined portion respectively.

We have also come across physical quantities such as displacement, velocity, acceleration, momentum etc, which are of different type in comparison to above. Consider the figure-1, where A, B, C are at a distance 4k.m. from P. If we start from P, then covering 4k.m. distance is not sufficient to describe the destination where we reach after the travel, the end point plays an important role giving rise the need of direction. So we need to study about direction of a quantity, along with magnitude.




Fig - 1

to  
So here  
of  
quantity,

### OBJECTIVE

After completion of the topic you are able to :-

- i) Define and distinguish between scalars and vectors.
- ii) Represent a vector as directed line segment.
- iii) Classify vectors in to different types.
- iv) Resolve vector along two or three mutually perpendicular axes.
- v) Define dot product of two vectors and explain its geometrical meaning.
- vi) Define cross product of two vectors and apply it to find area of triangle and parallelogram.

### Expected background knowledge

- i) Knowledge of plane and co-ordinate geometry
- ii) Trigonometry.

### Scalars and vectors

All the physical quantities can be divided into two types. i)

Scalar quantity or Scalar. ii) Vector quantity or Vector.

**Scalar quantity:** - The physical quantities which requires only magnitude for its complete specification is called as scalar quantities.

Examples: - Speed, mass, distance, velocity, volume etc.

**Vector:** - A directed line segment is called as vector.

**Vector quantities:-** A physical quantity which requires both magnitude & direction for its complete specification and satisfies the law of vector addition is called as vector quantities. Examples: - Displacement, force, acceleration, velocity, momentum etc. **Representation of vector:-** A vector is a directed line segment  $\overrightarrow{AB}$  where A is the initial point and B is the terminal point and direction is from A to B.

(see fig-2).

Similarly  $\overrightarrow{BA}$  is a directed line which represents a vector having initial point B and terminal point A.




Fig - 2

**Notation: -** A vector quantity is always represented by an arrow ( $\rightarrow$ ) mark over it or by bar ( $\overline{\phantom{a}}$ ) over it. For example  $\overrightarrow{AB}$ . It is also represented by a single small letter with an arrow or bar mark over it. For example  $\vec{a}$ .




Fig - 3

**Magnitude of a vector: -** Magnitude or modulus of a vector is the length of the vector. It is a scalar quantity.

Magnitude of  $\overrightarrow{AB} = |\overrightarrow{AB}| = \text{Length AB.} = AB$

**Types of Vector: -** Vectors are of following types.

- 1) Null vector or zero vector or void vector: -** A vector having zero magnitude and arbitrary direction is called as a null vector and is denoted by  $\vec{0}$ .

Clearly, a null vector has no definite direction. If  $\vec{a} = \overrightarrow{AB}$ , then  $\vec{a}$  is a null (or zero) vector iff  $|\vec{a}| = 0$  i.e. if

$$|\overrightarrow{AB}| = 0$$

For a null vector initial and terminal points are same.

- 2) Proper vector: -** Any non zero vector is called as a proper vector. If  $|\vec{a}| \neq 0$  then  $\vec{a}$  is a proper vector. **3) Unit vector: -** A vector whose magnitude is unity is called a unit vector. Unit vectors are denoted by a small letter with  $\hat{\phantom{a}}$  over it. For example  $\hat{a}$ .  $|\hat{a}| = 1$

5) **Co-Linear vectors:-** Vectors are said to be co-linear or parallel if they have the same line of action. In figure-5  $\vec{AB}$  and  $\vec{BC}$  are collinear.



Fig - 5

6) **Parallel vectors:** - Vectors are said to be parallel if they have same line of action or have line of action parallel to one another. In fig-6 the vectors are parallel to each other.




Fig - 6

7) **Co-planner Vectors:** - Vectors are said to be co-planner if they lie on the same plane. In fig-7 vector  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are coplanar.




Fig - 7

8) **Negative of a vector:** - A vector having same magnitude but opposite in direction to that of a given vector is called negative of that vector. If  $\vec{a}$  is any vector then negative vector of it is written as  $-\vec{a}$  and  $|\vec{a}| = |-\vec{a}|$  but both have direction opposite to each other as shown in fig-8.




Fig - 8

9) **Equal Vectors:** - Two vectors are said to be equal if they have same magnitude as well as same direction.




Fig - 9

Thus  $\vec{a} = \vec{b}$

Remarks:- Two vectors can not be equal

- i) If they have different magnitude .
- ii) If they have inclined supports. iii) If they have different sense.

**Vector operations Addition of vectors:** -

**Triangle law of vector addition:** - The law states that If two vectors are represented by the two sides of a triangle taken in same order their sum or resultant is represented by the 3<sup>rd</sup> side of the triangle with direction in reverse order.

As shown in figure-10  $\vec{a}$  and  $\vec{b}$  are two vectors represented by two sides OA and AB of a triangle

ABC in same order. Then the sum  $\vec{a} + \vec{b}$  is represented by the third side OB taken in reverse order i.e. the vector  $\vec{a}$  is represented by the directed segment  $\overrightarrow{OA}$  and the vector  $\vec{b}$  be the

directed segment  $\overrightarrow{AB}$ , so that the terminal point A of  $\vec{a}$  is

the initial point of  $\vec{b}$ . Then  $\overrightarrow{OB}$  represents the sum (or resultant)  $(\vec{a} + \vec{b})$ . Thus  $\overrightarrow{OB} = \vec{a} + \vec{b}$




Fig - 10

**Note-1** – The method of drawing a triangle in order to define the vector sum  $(\vec{a} + \vec{b})$  is called triangle law of addition of the vectors.

**Note-2** – Since any side of a triangle is less than the sum of the other two sides

$$|\overrightarrow{OB}| \neq |\overrightarrow{OA}| + |\overrightarrow{AB}|$$

**Parallelogram law of vector addition:** - If  $\vec{a}$  and  $\vec{b}$  are two vectors represented by two adjacent side of a parallelogram in magnitude and direction, then their sum (resultant) is represented in magnitude and direction by the diagonal which is passing through the common initial point of the two vectors.




Fig - 11

As shown in fig-II if OA is  $\vec{a}$  and AB is  $\vec{b}$  then OB diagonal represent  $\vec{a} + \vec{b}$  .

i.e.  $\vec{a} + \vec{b} = \overrightarrow{OA} + \overrightarrow{AB}$

**Polygon law of vector addition:** If  $\vec{a}, \vec{b}, \vec{c}$  and  $\vec{d}$  are the four sides of a polygon in same order then their sum is represented by the last side of the polygon taken in opposite order as shown in figure-12.




Fig - 12

**Subtraction of two vectors**

If  $\vec{a}$  and  $\vec{b}$  are two given vectors then the subtraction of  $\vec{b}$  from  $\vec{a}$  denoted by  $\vec{a} - \vec{b}$  is defined as addition of  $-\vec{b}$  with  $\vec{a}$ . i.e.  $\vec{a} - \vec{b} = \vec{a} + (-\vec{b})$ .

**Properties of vector addition:-** i) Vector addition is commutative i.e. if  $\vec{a}$  &  $\vec{b}$  are any two vectors then:-

$$\vec{a} + \vec{b} = \vec{b} + \vec{a}$$

ii) Vector addition is associative i.e. if  $\vec{a}, \vec{b}, \vec{c}$  are any three vectors,

$$\text{then } (\vec{a} + \vec{b}) + \vec{c} = \vec{a} + (\vec{b} + \vec{c})$$

- iii) Existence of additive identity i.e. for any vector  $\vec{a}$ ,  $\vec{0}$  is the additive identity i.e.  $\vec{0} + \vec{a} = \vec{a} + \vec{0} = \vec{a}$  where  $\vec{0}$  is a null vector.
- iv) Existence of additive Inverse :- If  $\vec{a}$  is any non zero vector then  $-\vec{a}$  is the additive inverse of  $\vec{a}$ , so that  $\vec{a} + (-\vec{a}) = (-\vec{a}) + \vec{a} = \vec{0}$

### Multiplication of a vector by a scalar :-

If  $\vec{a}$  is a vector and k is a nonzero scalar then the multiplication of the vector  $\vec{a}$  by the scalar k is a vector denoted by  $k\vec{a}$  whose magnitude  $|k|$  times that of  $\vec{a}$ . i.e  $k\vec{a} = |k| \times |\vec{a}| = k \times |\vec{a}|$  if  $k \geq 0$ .  
 $= (-k) \times |\vec{a}|$  if  $k < 0$ .

The direction of  $k\vec{a}$  is same as that of  $\vec{a}$  if k is positive and opposite as that of  $\vec{a}$  if k is negative.

$k\vec{a}$  and  $\vec{a}$  are always parallel to each other.

**Properties of scalar multiplication of vectors :-** If h and k

are scalars and  $\vec{a}$  and  $\vec{b}$  are given vectors then i)  $k(\vec{a} + \vec{b}) = k\vec{a} + k\vec{b}$

ii)  $(h+k)\vec{a} = h\vec{a} + k\vec{a}$ , (Distributive law) iii)

$(hk)\vec{a} = h(k\vec{a})$ , (Associative law) iv) 1.

$\vec{a} = \vec{a}$

v)  $0 \cdot \vec{a} = \vec{0}$

### Position Vector of a point

Let O be a fixed point called origin, let P be any other point, then the vector  $\vec{OP}$  is called position vector of the point P relative to O and is denoted by  $\vec{p}$ . As shown in figure-13, let AB be any vector, then applying triangle law of addition we have

$\vec{OA} + \vec{AB} = \vec{OB}$  where  $\vec{OA} = \vec{a}$  and  $\vec{OB} = \vec{b}$

$\Rightarrow \vec{AB} = \vec{OB} - \vec{OA} = \vec{b} - \vec{a}$

= (Position vector of B) - (Position vector of A)

**Section Formula:-** Let A and B be two points with position vector  $\vec{a}$  and  $\vec{b}$  respectively and P be a point on line segment AB, dividing it in the ratio m:n. internally. Then the position vector of P i.e.  $\vec{r}$  is

given by the formula:  $\vec{r} = \frac{m\vec{b} + n\vec{a}}{m+n}$

If P divides AB externally in the ratio m:n then  $\vec{r} =$

$$\frac{m\vec{b} - n\vec{a}}{m-n}$$

If P is the midpoint of AB then  $\vec{r} = \frac{\vec{a} + \vec{b}}{2}$




Fig - 13




Fig - 14



**Example-1 :-** Prove that by vector method the medians of a triangle are concurrent.

Solution:- Let ABC be a triangle where  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are the position vector of A,B and C respectively. We have to show that the medians of this triangle are concurrent.




Fig - 15

Let AD , BE and CF are the three medians of the triangle.

Now as D be the midpoint of BC, so position vector of D i.e.  $\vec{d} = \frac{\vec{b} + \vec{c}}{2}$  .

Let G be any point of the median AD which divides AD in the ratio 2:1 . Then position vector of G is given

$$\text{by } \vec{g} = \frac{2\vec{d} + \vec{a}}{2+1} = \frac{2\left(\frac{\vec{b} + \vec{c}}{2}\right) + 1\vec{a}}{3} \quad (\text{by applying section formula})$$

$$\Rightarrow \vec{g} = \frac{\vec{a} + \vec{b} + \vec{c}}{3}$$

Let G' be a point which divides BE in the ratio 2:1 ,

Position vector of E is  $\vec{e} = \frac{\vec{a} + \vec{c}}{2}$  .

Then position vector of G' is given by  $\vec{g}' = \frac{2\vec{e} + \vec{b}}{2+1} = \frac{2\left(\frac{\vec{a} + \vec{c}}{2}\right) + 1\vec{b}}{3}$  (by applying section formula)

$$\Rightarrow \vec{g}' = \frac{\vec{a} + \vec{b} + \vec{c}}{3}$$

As position vector of a point is unique , so  $G = G'$  .

Similarly if we take G'' be a point on CF dividing it in 2:1 ratio then the position vector of G'' will be same as that of G.

Hence G is the one point where three median meet.

∴ The three medians of a triangle are concurrent. (proved)

**Example2: -** Prove that i)  $|\vec{a} + \vec{b}| \leq |\vec{a}| + |\vec{b}|$  (It is known as Triangle Inequality).

ii)  $|\vec{a}| - |\vec{b}| \leq |\vec{a} - \vec{b}|$  iii)  $|\vec{a} - \vec{b}| \leq |\vec{a}| + |\vec{b}|$

### Components of vector in 2D

Let XOY be the co-ordinate plane and P(x,y) be any point in this plane.

The unit vector along direction of X axis i.e.  $\overrightarrow{OX}$  is denoted by  $\hat{i}$ .

The unit vector along direction of Y axis i.e.  $\overrightarrow{OY}$  is denoted by  $\hat{j}$ .

Then from figure-18 it is clear that  $\overrightarrow{OM} = x\hat{i}$  and  $\overrightarrow{ON} = y\hat{j}$ .

So, the position vector of P is given by

$$\vec{r} = x\hat{i} + y\hat{j}$$

And  $OP = |\overrightarrow{OP}| = r = \sqrt{x^2 + y^2}$




Fig-18

**Representation of vector in component form in 2D**

If  $\overrightarrow{AB}$  is any vector having end points A(  $x_1, y_1$ ) and B(  $x_2, y_2$ ) , then it can be represented by  $\overrightarrow{AB} = (x_2 - x_1)\hat{i} + (y_2 - y_1)\hat{j}$

**Components of vector in 3D**

Let P(x,y,z) be a point in space and  $\hat{i}$ ,  $\hat{j}$  and  $\hat{k}$  be the unit vectors along X axis , Y axis and Z axis respectively. (as shown in fig-19 ) Then the position vector of P is given by

$\overrightarrow{OP} = x\hat{i} + y\hat{j} + z\hat{k}$ , The vectors  $x\hat{i}$ ,  $y\hat{j}$ ,  $z\hat{k}$  are called the components of  $\overrightarrow{OP}$  along x-axis, y-axis and z-axis respectively. And  $OP = |\overrightarrow{OP}| = \sqrt{x^2 + y^2 + z^2}$




Fig-19

**Addition and scalar multiplication in terms of component form of vectors: -**

For any vector  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$  and  $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$

i)  $\vec{a} + \vec{b} = (a_1 + b_1)\hat{i} + (a_2 + b_2)\hat{j} + (a_3 + b_3)\hat{k}$  ii)

$\vec{a} - \vec{b} = (a_1 - b_1)\hat{i} + (a_2 - b_2)\hat{j} + (a_3 - b_3)\hat{k}$  iii) k

$\vec{a} = k a_1\hat{i} + k a_2\hat{j} + k a_3\hat{k}$ , where K is a scalar. iv)  $\vec{a} = \vec{b}$

$a_1\hat{i} + a_2\hat{j} + a_3\hat{k} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$

$a_1 = b_1, a_2 = b_2, a_3 = b_3$

**Representation of vector in component form in 3-D & Distance between two points:**

If  $\overrightarrow{AB}$  is any vector having end points A(  $x_1, y_1, z_1$ ) and B(  $x_2, y_2, z_2$ ) , then it can be represented by  $\overrightarrow{AB} =$

**Position vector of B – Position vector of A**

$$= (x_2\hat{i} + y_2\hat{j} + z_2\hat{k}) - (x_1\hat{i} + y_1\hat{j} + z_1\hat{k})$$

$$= (x_2 - x_1)\hat{i} + (y_2 - y_1)\hat{j} + (z_2 - z_1)\hat{k}$$

$$|\overline{AB}| = \sqrt{\quad}$$

**Example 3:-**

Show that the points A(2,6,3), B(1,2,7) and C(3,10,-1) are collinear.

Solution:- From given data Position vector of A,  $\overline{OA} = 2\hat{i} + 6\hat{j} + 3\hat{k}$ .

$$\text{Position vector of B, } \overline{OB} = \hat{i} + 2\hat{j} + 7\hat{k}$$

$$\text{Position vector of C, } \overline{OC} = 3\hat{i} + 10\hat{j} - \hat{k}$$

Now  $\overline{AB} = \overline{OB} - \overline{OA} = (1 - 2)\hat{i} + (2 - 6)\hat{j} + (7 - 3)\hat{k} = -\hat{i} - 4\hat{j} + 4\hat{k}$

$$\overline{AC} = \overline{OC} - \overline{OA} = (3 - 2)\hat{i} + (10 - 6)\hat{j} + (-1 - 3)\hat{k} = \hat{i} + 4\hat{j} - 4\hat{k}$$

$$= -(-\hat{i} - 4\hat{j} + 4\hat{k}) = -\overline{AB}$$

∴  $\overline{AB}$  or  $\overline{AC}$  are collinear.

∴ They have same support and common point A.

As 'A' is common to both vector, that proves A, B and C are collinear.

**Example-4:** - Prove that the points having position vector given by  $2\hat{i} - \hat{j} + \hat{k}$ ,  $\hat{i} - 3\hat{j} - 5\hat{k}$  and  $3\hat{i} - 4\hat{j} - 4\hat{k}$  form a right angled triangle. [2009(w)]

Solution :- Let A, B and C be the vertices of a triangle with position vectors  $2\hat{i} - \hat{j} + \hat{k}$ ,  $\hat{i} - 3\hat{j} - 5\hat{k}$  and  $3\hat{i} - 4\hat{j} - 4\hat{k}$  respectively

Then  $\overline{AB}$  = Position vector of B - Position vector of A.

$$= (1 - 2)\hat{i} + (-3 - (-1))\hat{j} + (-5 - 1)\hat{k} = -\hat{i} - 2\hat{j} - 6\hat{k}$$

$\overline{BC}$  = Position vector of C - Position vector of B.

$$= (3 - 1)\hat{i} + (-4 - (-3))\hat{j} + (-4 - (-5))\hat{k} = 2\hat{i} - \hat{j} + \hat{k}$$

$\overline{AC}$  = Position vector of C - Position vector of A.

$$= (3 - 2)\hat{i} + (-4 - (-1))\hat{j} + (-4 - 1)\hat{k} = \hat{i} - 3\hat{j} - 5\hat{k}$$

Now  $AB = |\overline{AB}| = \sqrt{(-1)^2 + (-2)^2 + (-6)^2} = \sqrt{1 + 4 + 36} = \sqrt{41}$

$BC = |\overline{BC}| = \sqrt{2^2 + (-1)^2 + 1^2} = \sqrt{4 + 1 + 1} = \sqrt{6}$

$AC = |\overline{AC}| = \sqrt{1^2 + (-3)^2 + (-5)^2} = \sqrt{1 + 9 + 25} = \sqrt{35}$

From above  $BC^2 + AC^2 = 6 + 35 = 41 = AB^2$ .

Hence ABC is a right angled triangle.

**Angle between the vectors:-**

As shown in figure-20 angle between two vectors  $\vec{RS}$  and  $\vec{PQ}$  can be determined as follows.

Let  $\vec{OB}$  be a vector parallel to  $\vec{RS}$  and  $\vec{OA}$  is a vector parallel to  $\vec{PQ}$  such that  $\vec{OB}$  and  $\vec{OA}$  intersect each other.

Then  $\theta = \angle AOB =$  angle between  $\vec{RS}$  and  $\vec{PQ}$ .

If  $\theta = 0$  then vectors are said to be parallel.

If  $\theta = \frac{\pi}{2}$  then vectors are said to be orthogonal or perpendicular.




Fig-20

**Dot Product or**

**Scalar product of vectors**

The scalar product of two vectors  $\vec{a}$  and  $\vec{b}$  whose magnitudes are,  $a$  and  $b$  respectively, where  $\theta$  is the angle between them is

$$\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos\theta = a b \cos\theta$$

**Scalar product of vectors**

scalar product of two vectors  $\vec{a}$  and  $\vec{b}$  whose magnitudes respectively denoted by  $a$  and  $b$  is defined as the scalar  $ab \cos\theta$  where  $\theta$  is the angle between  $\vec{a}$  and  $\vec{b}$  such that  $0 \leq \theta \leq \pi$ .

**Geometrical meaning of dot product**

$\vec{a}$  and  $\vec{b}$  are two vectors having

In figure 21(a),

$\theta$  is the angle between  $\vec{a}$  and  $\vec{b}$ .

$\vec{b}$  on  $\vec{a}$  and from

angle between them is clear that,

Let  $M$  be the foot of perpendicular from  $B$  to  $OA$ .

Let  $OM$  be the projection of  $\vec{b}$  on  $\vec{a}$ .

perpendicular projection of  $\vec{b}$  on  $\vec{a} = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|}$

drawn from  $B$  to  $OA$ .  $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos\theta = |\vec{b}| |\vec{a}| \cos\theta$

Then  $OM$  is

$$|\vec{b}| \cos\theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|}$$

which gives

Similarly we can write

$$|\vec{a}| \cos\theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$$




Fig-21

the Projection of figure-21(a)

$$|\vec{a}| \cos\theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$$

$\vec{b} \cdot \vec{a}$  Similarly, let us draw a perpendicular from A on OB and let N be the foot of the perpendicular in fig-21(b).

Then  $ON = \text{Projection of } \vec{a} \text{ on } \vec{b}$   
 and  $ON = OA \cos \theta = |\vec{a}| \cos \theta$




Fig-21(b)

**Properties of Dot product**

- i)  $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{a}$  (commutative)
- ii)  $\vec{a} \cdot (\vec{b} + \vec{c}) = \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c}$  (Distributive)
- iii) If  $\vec{a} \parallel \vec{b}$ , then  $\vec{a} \cdot \vec{b} = ab$  { as  $\theta = 0$  in this case  $\cos 0 = 1$  }  
 In particular  $(\vec{a})^2 = \vec{a} \cdot \vec{a} = |\vec{a}|^2$   
 $\hat{i} \cdot \hat{i} = \hat{j} \cdot \hat{j} = \hat{k} \cdot \hat{k} = 1$
- iv) If  $\vec{a} \perp \vec{b}$ , then  $\vec{a} \cdot \vec{b} = 0$ . { as  $\theta = 90^\circ$  in this case  $\cos 90^\circ = 0$  }

In particular  $\hat{i} \cdot \hat{j} = \hat{j} \cdot \hat{k} = \hat{k} \cdot \hat{i} = 0 = \hat{j} \cdot \hat{i} = \hat{k} \cdot \hat{j} = \hat{i} \cdot \hat{k}$  v)  $\vec{a} \cdot \vec{0} = \vec{0} \cdot \vec{a} = 0$   
 vi)  $(\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b}) = |\vec{a}|^2 - |\vec{b}|^2 = a^2 - b^2$  {Where  $|\vec{a}| = a$  and  $|\vec{b}| = b$  }

viii) Work done by a Force:- The work done by a force  $\vec{F}$  acting on a body causing displacement  $\vec{d}$  is given by  $W = \vec{F} \cdot \vec{d}$

**Dot product in terms of rectangular components**

For any vectors  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$  and  $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$  we have,  
 $\vec{a} \cdot \vec{b} = a_1b_1 + a_2b_2 + a_3b_3$  ( by applying distributive( ii ), (iii) and (iv) successively)

**Angle between two non zero vectors**

For any two non zero vectors  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$  and  $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$ , having  $\theta$  is the angle between them we have,

$$\cos \theta = \frac{\vec{a} \cdot \vec{b}}{ab} = \hat{a} \cdot \hat{b} = \frac{a_1b_1 + a_2b_2 + a_3b_3}{\sqrt{a_1^2 + a_2^2 + a_3^2} \sqrt{b_1^2 + b_2^2 + b_3^2}} \text{ (In terms of components.)}$$

$$\theta = \cos^{-1} \left( \frac{a_1b_1 + a_2b_2 + a_3b_3}{\sqrt{a_1^2 + a_2^2 + a_3^2} \sqrt{b_1^2 + b_2^2 + b_3^2}} \right)$$

**Condition of Perpendicularity: -**

Two vectors  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$  and  $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$  are perpendicular to each other

$$\square a_1b_1 + a_2b_2 + a_3b_3 = 0$$

**Condition of Parallelism :-**

Two vectors  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$  and  $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$  are parallel to each other  $\square \frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3} = \dots$

**Scalar & vector projections of two vectors (Important formulae)**

Scalar Projection of  $\vec{b}$  on  $\vec{a} = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|}$

Vector Projection of  $\vec{b}$  on  $\vec{a} = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|^2} \hat{a} = \left[ \frac{\vec{b} \cdot \vec{a}}{|\vec{a}|^2} \right] \vec{a}$

Scalar Projection of  $\vec{a}$  on  $\vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$

Vector Projection of  $\vec{a}$  on  $\vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|^2} \hat{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|^2} \vec{b}$

**Examples: -**

**Q.- 7.** Find the value of p for which the vectors  $3\hat{i} + 2\hat{j} + 9\hat{k}$ ,  $\hat{i} + p\hat{j} + 3\hat{k}$  are perpendicular to each other.

Solution:- Let  $\vec{a} = 3\hat{i} + 2\hat{j} + 9\hat{k}$  and  $\vec{b} = \hat{i} + p\hat{j} + 3\hat{k}$ .

Here  $a_1 = 3, a_2 = 2, a_3 = 9$   $b_1 = 1, b_2 = p$  &  $b_3 = 3$

Given  $\vec{a} \perp \vec{b} \Rightarrow a_1b_1 + a_2b_2 + a_3b_3 = 0$

$$\Rightarrow 3 \cdot 1 + 2 \cdot p + 9 \cdot 3 = 0$$

$$\Rightarrow 3 + 2p + 27 = 0$$

$$\Rightarrow 2p = -30$$

$$\Rightarrow p = -15 \quad (\text{Ans})$$

**Q-8** Find the value of p for which the vectors  $\vec{a} = 3\hat{i} + 2\hat{j} + 9\hat{k}$ ,  $\vec{b} = \hat{i} + p\hat{j} + 3\hat{k}$  are parallel to each other.

( 2014-W)

Solution:- Given  $\vec{a} \parallel \vec{b} \Leftrightarrow \frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3} \Leftrightarrow \frac{3}{1} = \frac{2}{p} = \frac{9}{3}$  { Taking 1<sup>st</sup> two terms }

$$\square 3 = \frac{2}{p} \square p = \frac{2}{3} \quad (\text{Ans}) \quad \{\text{Note: - any two expression may be taken for finding p.}\}$$

**Q-9** Find the scalar product of  $3\hat{i} - 4\hat{j}$  and  $-2\hat{i} + \hat{j}$ . (2015-S) Solution:- (

$$3\hat{i} - 4\hat{j}) \cdot (-2\hat{i} + \hat{j}) = (3 \times (-2)) + ((-4) \times 1) = (-6) + (-4) = -10$$

**Q-10** Find the angle between the vectors  $5\hat{i} + 3\hat{j} + 4\hat{k}$  and  $6\hat{i} - 8\hat{j} - \hat{k}$ . (2015-W)

Solution:- Let  $\vec{a} = 5\hat{i} + 3\hat{j} + 4\hat{k}$  and  $\vec{b} = 6\hat{i} - 8\hat{j} - \hat{k}$  Let  $\theta$  be the angle between  $\vec{a}$  and  $\vec{b}$ .

$$\theta = \cos^{-1}\left(\frac{a_1b_1 + a_2b_2 + a_3b_3}{\sqrt{a_1^2 + a_2^2 + a_3^2}\sqrt{b_1^2 + b_2^2 + b_3^2}}\right)$$

Then

$$= \cos^{-1}\left(\frac{5.6 + 3.(-8) + 4.(-1)}{\sqrt{5^2 + 3^2 + 4^2}\sqrt{6^2 + (-8)^2 + (-1)^2}}\right) = \cos^{-1}\left(\frac{30 - 24 - 4}{\sqrt{50}\sqrt{101}}\right) = \cos^{-1}\left(\frac{2}{\sqrt{50}\sqrt{101}}\right)$$

**Q-11** Find the scalar and vector projection of  $\vec{a}$  on  $\vec{b}$  where,  
 $\vec{a} = \hat{i} - \hat{j} - \hat{k}$  and  $\vec{b} = 3\hat{i} + \hat{j} + 3\hat{k}$ . { 2013-W, 2017-W, 2017-S }

Solution:- Scalar Projection of  $\vec{a}$  on  $\vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} = \frac{1.3 + (-1).1 + (-1).3}{\sqrt{3^2 + 1^2 + 3^2}} = \frac{3 - 1 - 3}{\sqrt{19}} = \frac{-1}{\sqrt{19}}$

$$\text{Vector Projection of } \vec{a} \text{ on } \vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|^2} \vec{b} = \frac{1.3 + (-1).1 + (-1).3}{(\sqrt{3^2 + 1^2 + 3^2})^2} (3\hat{i} + \hat{j} + 3\hat{k})$$

$$= \frac{3 - 1 - 3}{19} (3\hat{i} + \hat{j} + 3\hat{k}) = \frac{-1}{19} (3\hat{i} + \hat{j} + 3\hat{k})$$

**Q-12** Find the scalar and vector projection of  $\vec{b}$  on  $\vec{a}$  where,  
 $\vec{a} = 3\hat{i} + \hat{j} - 2\hat{k}$  and  $\vec{b} = 2\hat{i} + 3\hat{j} - 4\hat{k}$ . { 2015-S }

Solution: - Scalar Projection of  $\vec{b}$  on  $\vec{a} = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|} = \frac{3.2 + 1.3 + (-2).(-4)}{\sqrt{3^2 + 1^2 + (-2)^2}} = \frac{6 + 3 + 8}{\sqrt{14}} = \frac{17}{\sqrt{14}}$

$$\text{Vector Projection of } \vec{b} \text{ on } \vec{a} = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|^2} \vec{a} = \frac{3.2 + 1.3 + (-2).(-4)}{(\sqrt{3^2 + 1^2 + (-2)^2})^2} (3\hat{i} + \hat{j} - 2\hat{k})$$

$$= \frac{17}{14} (3\hat{i} + \hat{j} - 2\hat{k})$$

**Q-13** If  $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c}$ , then prove that  $\vec{a} = \vec{0}$  or  $\vec{b} = \vec{c}$  or  $\vec{a} \perp (\vec{b} - \vec{c})$

Proof:- Given  $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c}$

$$\Rightarrow (\vec{a} \cdot \vec{b}) - (\vec{a} \cdot \vec{c}) = 0 \Rightarrow \vec{a} \cdot (\vec{b} - \vec{c}) = 0 \quad \{ \text{applying distributive property} \}$$

Dot product of above two vector is zero indicates the following conditions  $\vec{a} = \vec{0}$   
or  $\vec{b} - \vec{c} = \vec{0}$  or  $\vec{a} \perp (\vec{b} - \vec{c})$

$$\Rightarrow \vec{a} = \vec{0} \text{ or } \vec{b} = \vec{c} \text{ or } \vec{a} \perp (\vec{b} - \vec{c}) \quad (\text{proved})$$

**Example:-14** Find the work done by the force  $\vec{F} = \hat{i} + \hat{j} - \hat{k}$  acting on a particle if the particle is displaced A from A(3,3,3) to B(4,4,4). Ans:- Let O be the origin, then

$$\text{Position vector of A } \vec{OA} = 3\hat{i} + 3\hat{j} + 3\hat{k}$$

$$\text{Position vector of B } \vec{OB} = 4\hat{i} + 4\hat{j} + 4\hat{k}$$

Then displacement is given by,  $\vec{d} = \vec{AB} = (\vec{OB} - \vec{OA}) = (4\hat{i} + 4\hat{j} + 4\hat{k}) - (3\hat{i} + 3\hat{j} + 3\hat{k}) = \hat{i} + \hat{j} + \hat{k}$

So work done by the force  $W = \vec{F} \cdot \vec{d} = \vec{F} \cdot \vec{AB} = (\hat{i} + \hat{j} - \hat{k}) \cdot (\hat{i} + \hat{j} + \hat{k})$

$$= 1.1 + 1.1 + (-1).1 = 1 \text{ units}$$

**Example:-15** If  $\hat{a}$  and  $\hat{b}$  are two unit vectors and  $\theta$  is the angle between them then prove that

$$\sin \frac{\theta}{2} = \frac{1}{2} |\hat{a} - \hat{b}|$$

Proof: -  $(|\hat{a} - \hat{b}|)^2 = (\hat{a} - \hat{b}) \cdot (\hat{a} - \hat{b}) = (\hat{a} \cdot \hat{a}) - (\hat{a} \cdot \hat{b}) - (\hat{b} \cdot \hat{a}) + (\hat{b} \cdot \hat{b})$  { Distributive property }

$$= (|\hat{a}|)^2 - (\hat{a} \cdot \hat{b}) - (\hat{a} \cdot \hat{b}) + (|\hat{b}|)^2$$
 { commutative property }

$$= 1^2 - 2 \hat{a} \cdot \hat{b} + 1^2$$
 { as  $\hat{a}$  and  $\hat{b}$  are unit vectors so their magnitudes are 1 }

$$= 2 - 2 \hat{a} \cdot \hat{b} = 2 (1 - \hat{a} \cdot \hat{b})$$

$$= 2 (1 - |\hat{a}| |\hat{b}| \cos \theta)$$
 { as  $\theta$  is the angle between  $\hat{a}$  and  $\hat{b}$  }

$$= 2 (1 - 1 \cdot 1 \cdot \cos \theta)$$

$$= 2 (1 - \cos \theta) = 2 \cdot 2 \sin^2 \frac{\theta}{2}$$

Taking square root of both sides we have  $|\hat{a} - \hat{b}| = 2 \sin \frac{\theta}{2}$

$$\square \sin \frac{\theta}{2} = \frac{1}{2} |\hat{a} - \hat{b}| \text{ (proved)}$$

**Example:-16** If the sum of two unit vectors is a unit vector. Then show that the magnitude of their difference is  $\sqrt{3}$ .

Proof:-  $\hat{a}, \hat{b}$  and  $\hat{c}$  are three unit vectors such that  $\hat{a} + \hat{b} = \hat{c}$

Squaring both sides we have,  $\Rightarrow$

$$(|\hat{a} + \hat{b}|)^2 = |\hat{c}|^2 = 1$$

$$(|\hat{a}|)^2 + (|\hat{b}|)^2 + 2 \hat{a} \cdot \hat{b} = 1$$

$$\Rightarrow 1 + 1 + 2 \hat{a} \cdot \hat{b} = 1$$

$$\Rightarrow 1^2 + 1^2 + 2 |\hat{a}| |\hat{b}| \cos \theta = 1$$
 { where  $\theta$  is the angle between  $\hat{a}$  and  $\hat{b}$  }

$$\Rightarrow 1 + 1 + 2 \cos \theta = 1$$

$$\Rightarrow 2 \cos \theta = -1$$

$$\Rightarrow \cos \theta = \frac{-1}{2}$$

Now we have to find the magnitude of their difference i.e  $|\hat{a} - \hat{b}|$ .

$$\text{So } (|\hat{a} - \hat{b}|)^2 = (|\hat{a}|)^2 + (|\hat{b}|)^2 - 2 \hat{a} \cdot \hat{b} = 1^2 + 1^2 - 2 |\hat{a}| |\hat{b}| \cos \theta$$

$$= 2 - 2 \cos \theta = 2 - 2 \left(\frac{-1}{2}\right) = 2 - (-1) = 3$$

$$\therefore |\hat{a} - \hat{b}| = \sqrt{3} \text{ (Proved)}$$



### Vector Product or Cross Product

If  $\vec{a}$  and  $\vec{b}$  are two vectors and  $\theta$  is the angle between them, then the vector product of these two vectors denoted by  $\vec{a} \times \vec{b}$  is defined as

$$\vec{a} \times \vec{b} = |\vec{a}| |\vec{b}| \sin \theta \hat{n}$$

where  $\hat{n}$  is the unit vector perpendicular to both  $\vec{a}$  and  $\vec{b}$ . As shown in figure-21 the direction of  $\vec{a} \times \vec{b}$  is always perpendicular to both  $\vec{a}$  and  $\vec{b}$ .




Fig-22

### Properties of cross product

- i) Vector product is not commutative  $\vec{a} \times \vec{b} \neq \vec{b} \times \vec{a}$
- ii) For any two vectors  $\vec{a}$  and  $\vec{b}$ ,  $\vec{a} \times \vec{b} = -(\vec{b} \times \vec{a})$
- iii) For any scalar  $m$ ,  $m(\vec{a} \times \vec{b}) = (m\vec{a}) \times \vec{b} = \vec{a} \times (m\vec{b})$
- iv) Distributive  $\vec{a} \times (\vec{b} + \vec{c}) = (\vec{a} \times \vec{b}) + (\vec{a} \times \vec{c})$
- v) Vector product of two parallel or collinear vectors is zero.

$$\vec{a} \times \vec{a} = \vec{0} \quad \text{and if } \vec{a} \parallel \vec{b} \text{ then } \vec{a} \times \vec{b} = \vec{0} \quad \{ \text{as } \theta = 0 \text{ or } 180^\circ \Rightarrow \sin \theta = 0 \}$$

Using this property we have,

$$\hat{i} \times \hat{i} = \hat{j} \times \hat{j} = \hat{k} \times \hat{k} = \vec{0}$$

v) Vector product of orthonormal unit vectors form a right handed system. As shown in figure- 23 the three mutually perpendicular unit vectors  $\hat{i}, \hat{j}, \hat{k}$  form a right handed system, i.e.  $\hat{i} \times \hat{j} = \hat{k} = -(\hat{j} \times \hat{i})$

( as  $\theta=90$  , then  $\sin\theta = 1$ )

$$\hat{j} \times \hat{k} = \hat{i} = -(\hat{k} \times \hat{j})$$

$$\hat{k} \times \hat{i} = \hat{j} = -(\hat{i} \times \hat{k})$$




Fig-23

**Unit vector perpendicular to two vectors:-** Unit vector perpendicular to two given vectors  $\vec{a}$  and  $\vec{b}$  is

given by 
$$\hat{n} = \frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|}$$

**Angle between two vectors**

Let  $\theta$  be the angle between  $\vec{a}$  and  $\vec{b}$ . Then  $\vec{a} \times \vec{b} = (|\vec{a}| |\vec{b}| \sin\theta) \hat{n}$ .

Taking modulus of both sides

we have,  $|\vec{a} \times \vec{b}| = |\vec{a}| |\vec{b}| \sin\theta$

$|\vec{a} \times \vec{b}| = |\vec{a}| |\vec{b}| \sin\theta$

$$\sin\theta = \frac{|\vec{a} \times \vec{b}|}{|\vec{a}| |\vec{b}|}$$

Hence 
$$\theta = \sin^{-1} \left\{ \frac{|\vec{a} \times \vec{b}|}{|\vec{a}| |\vec{b}|} \right\}$$

**Geometrical Interpretation of vector product or cross product**

Let  $\vec{OA} = \vec{a}$  and  $\vec{OB} = \vec{b}$ .

Then  $\vec{a} \times \vec{b} = (|\vec{a}| |\vec{b}| \sin\theta) \hat{n}$

$$= (|\vec{a}| |\vec{b}| \sin\theta) \hat{n}$$

From fig-24 below it is clear that

$$BM = OB \sin\theta = |\vec{b}| \sin\theta = |\vec{a}| |\vec{b}| \sin\theta \hat{n}$$

{ as  $\sin\theta = BM/OB$  &  $\vec{OB} = \vec{b}$ }

Now  $|\vec{a} \times \vec{b}| = |\vec{a}| |\vec{b}| \sin\theta = |\vec{a}| BM$

$BM =$  Area of the parallelogram with side  $\vec{a}$  and  $\vec{b}$ .




Fig-24

Therefore the magnitude of cross product of two vectors is equal to area of the parallelogram formed by these vectors as two adjacent sides.

From this it can be concluded that area of  $\Delta ABC = \frac{1}{2} \overline{IAB} \cdot \overline{ACI}$

**Application of cross product**

1. Moment of a force about a point ( $\vec{M}$ ) :-Let O be any point and Let  $\vec{r}$  be the position vector w.r.t. O of

any point 'P' on the line of action of the force  $F$ , then the moment or torque of the force F about origin 'O' is given by

$$\vec{M} = \vec{r} \times \vec{F}$$

2. If  $\vec{a}$  and  $\vec{b}$  represent two adjacent sides of a triangle then the area of the triangle is given by  $\Delta = \frac{1}{2} |\vec{a} \times \vec{b}|$  Sq. unit

3. If  $\vec{a}$  and  $\vec{b}$  represent two adjacent sides of a parallelogram then area of the parallelogram is given by  $\Delta = |\vec{a} \times \vec{b}|$  Sq. unit

4. If  $\vec{a}$  and  $\vec{b}$  represent two diagonals of a parallelogram then area of the parallelogram is given by  $\Delta = \frac{1}{2} |\vec{a} \times \vec{b}|$  Sq. unit

**Vector product in component form :-**

If  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$  and  $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$ .

$$\begin{aligned} \vec{a} \times \vec{b} &= (a_1\hat{i} + a_2\hat{j} + a_3\hat{k}) \times (b_1\hat{i} + b_2\hat{j} + b_3\hat{k}) \\ &= a_1b_1(\hat{i} \times \hat{i}) + a_1b_2(\hat{i} \times \hat{j}) + a_1b_3(\hat{i} \times \hat{k}) + a_2b_1(\hat{j} \times \hat{i}) + a_2b_2(\hat{j} \times \hat{j}) + a_2b_3(\hat{j} \times \hat{k}) \\ &\quad + a_3b_1(\hat{k} \times \hat{i}) + a_3b_2(\hat{k} \times \hat{j}) + a_3b_3(\hat{k} \times \hat{k}) \end{aligned}$$

{ using properties  $\hat{i} \times \hat{i} = \hat{j} \times \hat{j} = \hat{k} \times \hat{k} = \vec{0}$ ,  $\hat{i} \times \hat{j} = \hat{k} = -(\hat{j} \times \hat{i})$ ,  $\hat{j} \times \hat{k} = \hat{i} = -(\hat{k} \times \hat{j})$  and  $\hat{k} \times \hat{i} = \hat{j} = -(\hat{i} \times \hat{k})$  }

$$\begin{aligned} &= (a_2b_3 - a_3b_2)\hat{i} + (a_3b_1 - a_1b_3)\hat{j} + (a_1b_2 - a_2b_1)\hat{k} \\ &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} \quad \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} \end{aligned}$$

i.e.

**Condition of Co-planarity**

If three vectors  $\vec{a}, \vec{b}$  and  $\vec{c}$  lies on the same plane then the perpendicular to  $\vec{a}$  and  $\vec{b}$  must be perpendicular to  $\vec{c}$ .

In particular  $(\vec{a} \times \vec{b}) \perp \vec{c} \Rightarrow (\vec{a} \times \vec{b}) \cdot \vec{c} = 0$

In component form if  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}, \vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$  and  $\vec{c} = c_1\hat{i} + c_2\hat{j} + c_3\hat{k}$

Then  $(\vec{a} \times \vec{b}) \cdot \vec{c} = 0$

## UNIT-II

# LIMITS AND CONTINUITY

### INTRODUCTION:-

The concept of limit and continuity is fundamental in the study of calculus. The fragments of this concept are evident in the method of exhaustion formulated by ancient Greeks and used by Archimedes (287-212 BC) in obtaining a formula for the area of the circular region conceived as successive approximation of areas of inscribed polygons with increased number of sides.

The concept of calculus is used in many engineering fields like Newton's Law of cooling derivation of basic Fluid mechanics equation etc.

In general the study of the theory of calculus mainly depends upon functions. Thus it is desirable to discuss the idea of functions before study of calculus.

### OBJECTIVES:-

- After studying this topic, you will
- be able to (i) Define function and cited examples there of
  - (ii) State types of functions.
  - (iii) Define limit of a function.
  - (iv) Evaluate limit of a function using different methods.
  - (v) Define continuity of a function at a point.
  - (vi) Test continuity of a function at a point.

### EXPECTED BACKGROUND KNOWELDGE :-

- (1) Set Theory
- (2) Concept of order pairs.

### RELATION:-

**Definition :** - If A and B are two nonempty sets, then any subset of  $A \times B$  is called a relation 'R' from A to B.

Mathematically  $R \subset A \times B$

Since  $\emptyset \subset A \times B$  and  $A \times B$  is a subset of itself, therefore  $\emptyset$  and  $A \times B$  are relations from A to B.

EXAMPLE:-1

$$A = \{1,2\} \quad B = \{a, b, c\}$$

Then  $A \times B = \{(1,a), (1,b), (1,c), (2,a), (2,b), (2,c)\}$

Now the following subsets of  $A \times B$  give some examples of relation from A to B.

$R_1 = \{(1, a), (2, b)\}$ ,  $R_2 = \{(1, c)\}$ ,  $R_3 = \{(2, b)\}$  and  $R_4 = \{(1, a), (1, b), (2, c)\}$  are examples of some relations.

The above relations can be represented by figure1-4 as follows.

$R_1 = \{(1, a), (2, b)\}$




Fig-1

$R_2 = \{(1, c)\}$




Fig-2

$R_3 = \{(2, b)\}$




Fig-3

$R_4 = \{(1, a), (1, b), (2, c)\}$




Fig-4

**Example – 2**

$A = \text{Players} = \{sachin, dhoni, pele, saina\}$

$B = \text{Game} = \{cricket, badminton, football\}$

Then  $R = \text{player related to their games.}$

The pictorial representation is given in Fig-5




Fig-5

**FUNCTION: -**

A relation ‘f’ from X to Y is called a function if it satisfies the following two conditions (i)

All elements of X are related to the elements of Y.

(ii) Each element of X related to only one element of Y.

In above example -1 only relation  $R_1$  is a function, because all elements of A is related.

Each element (i.e. 1-a and 2-b) is related to only one element of B.

$R_2$  is not a function as ‘2’ is not related.

$R_3$  is not a function as ‘1’ is not related.

$R_4$  is not a function as 1 related a and b.

EXAMPLE – 2 represent a function.

**EXAMPLE -4**

Let us consider a function F from  $A = \{1,2,3\}$  to  $B = \{a, b, c, d\}$  as follows.




Fig-6

Here we can write  $F(1) = a$

$F(2) = c$

$F(3) = b$

**DOMAIN ( $D_F$ )**

Let  $F: X \rightarrow Y$  is a function then the First set ‘X’ is called domain of F.

$$X = \text{Dom } F = D_F$$

In example-4  $\{1,2,3\}$  is the domain.

**Co-domain –**

If  $F: X \rightarrow Y$  is a function, then the 2<sup>nd</sup> set  $Y$  is called co-domain of  $F$ . In example -4  $\{a, b, c, d\}$  is the codomain.

**IMAGE:-** If  $f: X \rightarrow Y$  is a function and for any  $x \in X$ , we have  $f(x) \in Y$ .

This  $f(x) = y \in Y$  is called the image of  $x$ .

In example -4  
                   ‘a’ is the image of 1  
                   ‘b’ is the image of 3  
                   ‘c’ is the image of 2

**Range ( $R_F$ ): -**

The image set of ‘ $X$ ’ i.e. domain is called range of  $F$ .

$$F(X) = \text{Range of } F$$

In example -4  $\{a, b, c\}$  represent the range of  $F$ .

In above discussion we have taken examples of finite sets. But when we consider infinite sets it is not possible to represent a function either in tabular form or in figure form. So, we define function in another way as follows.

**CONSTANT-** A quantity which never changes its value. Constants are denoted by  $A, B, C$  etc.

**VARIABLE:-** A quantity which changes its value continuously  $x, y, z$  etc are used for variables.

**TWO TYPES OF VARIABLE:-** i) Independent variable ii) dependent variable

Independent variable  $\square$  Variable which changes its value independently. Generally we take ‘ $x$ ’ as independent variable.

Dependent Variable  $\square$  Variable which changes its value depending upon independent variable. We take ‘ $y$ ’ as dependent Variable.

**DEFINITION OF FUNCTION:-**

Let  $X$  and  $Y$  be two non empty sets. Then a function or mapping ‘ $f$ ’ assigned from set  $X$  to the set  $Y$  is a sort of correspondence which associated to each element  $x \in X$  a unique element  $y \in Y$  and is written as  $f: X \rightarrow Y$  ( read as “  $f$  maps  $X$  into  $Y$  ).

The element 'y' is called the image of x under f and is denoted by f(x) i.e.  $y = f(x)$  and x is called pre-image of y.

**Example -** Let  $y = F(x) = x^2$  where  $X = \{ 1, 1.5, 2 \}$

Pictorial representation is given in Fig-7




Fig-7

Here values of x form domain and values of y form range.

**FUNCTIONAL VALUE f(a)** □ The value of f(x) obtained by replacing x by a is called functional value of f(x) at x=a, denoted by f(a)

Example :- Let  $y = f(x) = x^2$

Then functional value of f(x) at x=2 is  $f(2) = 2^2 = 4$

The functional value of f(x) at x=1.5 is  $f(1.5) = 2.25$   
 If  $f(x) = \frac{1}{x}$  then  $f(1) = 1/1 = 1$ ,  $f(2) = 1/2$ . But  $f(0) = \frac{1}{0}$  which is undefined

So the function value of function is either finite or undefined.

**Classification of function**

Functions are classified into following categories

**Into function**

A function  $F : A \rightarrow B$  is said to be into if there exist at least one element in B which has no pre-image in A.

In this case Range set is a proper subset of co-domain Y.

Let  $A = \{ 1, 2, 3 \}$  and  $B = \{ a, b, c, d \}$

Then the function F given by fig-8 represent one into function from A to B.




Fig-8 ( Into Map)

In above figure d has no pre-image in A.



**Onto function**

A function  $F : A \rightarrow B$  is said to be onto if Range of F i.e.  $F(A) = B$ . In other words every element of B has a pre-image in A.

Let  $A = \{1,2,3,4\}$  and  $B = \{a,b,c\}$

Then the function F given by fig-9 represent one onto function from X to Y.




fig-9  
(Onto Map)


From above figure it is clear that  $F(A) = B$ .

**One-one function**

A function  $F: A \rightarrow B$  is said to be one-one if each distinct elements in A have distinct images in B i.e. if  $x_1 \neq x_2$  in A  $\Rightarrow F(x_1) \neq F(x_2)$  in B.

Let  $A = \{a,b,c\}$  and  $B = \{1,2,3,4\}$

Then the function F given by fig-10 represent an one-one function from A to B .



one - one


Fig-10

**Many-one function**

A function  $F : A \rightarrow B$  is said to be many-one if there exists at least one element in B, which has more than one pre-image in A.

Let  $A = \{1,2,3\}$  and  $B = \{a,b\}$

Then the function F given by fig-11 represent a many-one function from A to B .




( Many one)  
Fig-11

From above figure it is clear that a has two pre-images 1 and 2 in A .

**One-one and onto function or bijective function** A function  $F : A \rightarrow B$  is said to be one-one and onto if it is one-one and onto i.e each distinct element in A has distinct image in B and every element of B has a pre-image in A.

Let  $A = \{1,2,3\}$  and  $B = \{a,b,c\}$

Then the function F given by fig-12 represent an one-one and onto function from A to B .



( One-one and onto)

Fig-12

**Inverse function:-**

If  $F : X \rightarrow Y$  is a bijective function then it's inverse function defined from Y to X denoted by  $F^{-1}$  . If  $x \in X$  and  $y \in Y$  such that  $y = F(x)$  then  $x = F^{-1}(y)$ .

One example of inverse function is given in Fig-13 .

Let  $X = \{a,b,c\}$  and  $Y = \{1,2,3\}$  .




Fig-13

**Composition of two function**

If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  be any two functions having Range f = Domain g, then composition of f and g denoted by  $g \circ f$  is defined by  $g \circ f(x) = g(f(x))$ ,  $x \in X$ .

Domain of  $g \circ f = X$  and Range of  $g \circ f =$  Range of g. The composition of two function f and g is shown in fig-14.




Fig-14

Examples of some composite functions is given below

$y = f(x) = \sin x^2$  formed by composition of  $x^2$  and  $\sin x$ .

$y = f(x) = \sqrt{\cot x}$  formed by composition of  $\cot x$  and  $\sqrt{x}$  .

$y = f(x) = \log_a \sin \sqrt{x}$  formed by composition of  $\sqrt{x}$  ,  $\sin x$  and  $\log_a x$

**Real Valued Function:-**

$F : X \rightarrow Y$  is called a real valued Function.

If  $\text{dom } F = X \subset \mathbb{R}$  and  $Y \subset \mathbb{R}$ .

Generally we discuss our topics on this type of functions.

**DIFFERENT TYPES OF FUNCTIONS:-**

**(1) CONSTANT FUNCTION :-**

The function  $F(x) = K$  for all  $x \in R$ , where  $K$  is some real number is called a constant function

For Constant Function  $D_f = R$

$$R_f = \{K\}$$

**(2) IDENTITY FUNCTION :-**

$F(x) = x \forall x \in R$ , is called Identity Function. ( $\forall x$  means for all  $x$ ) It is also denoted by  $I_x$  or  $I$ .

$Dom_f = D_f = R$
-------------------

$$R = R$$

**(3) TRIGONOMETRIC FUNCTIONS :-**

$\sin x, \cos x, \tan x, \cot x, \sec x, \operatorname{cosec} x$  are called trigonometric functions. we know the definition of these functions.

We know that  $-1 \leq \sin x \leq 1$  and  $-1 \leq \cos x \leq 1 \quad \forall x \in R$ . Here  $x$  is radian measure of an angle.

Function	Domain	Range
Sinx	R	[-1,1]
Cosx	R	[-1,1]
Tanx	$R - \{(2n+1)\frac{\pi}{2} : n \in Z\}$	R
Cotx	$R - \{n\pi : n \in Z\}$	R
Secx	$R - \{(2n+1)\frac{\pi}{2} : n \in Z\}$	$R - (-1,1)$
Cosecx	$R - \{n\pi : n \in Z\}$	$R - (-1,1)$

**(4) INVERSE TRIGONOMETRIC FUNCTIONS :-**

$\sin^{-1}x, \cos^{-1}x, \tan^{-1}x, \cot^{-1}x, \sec^{-1}x, \operatorname{cosec}^{-1}x$  are called inverse trigonometric functions.

These are real functions.

Function	Domain	Range
$\sin^{-1}x$	[-1,1]	$[-\frac{\pi}{2}, \frac{\pi}{2}]$
$\cos^{-1}x$	[-1,1]	$[0, \pi]$

$\tan^{-1}x$	R	$(-\frac{\pi}{2}, \frac{\pi}{2})$
$\cot^{-1}x$	R	$(0, \pi)$
$\sec^{-1}x$	$x \leq -1$ or $x \geq 1$	$0 \leq y \leq \pi, y \neq \frac{\pi}{2}$
$\operatorname{cosec}^{-1}x$	$x \leq -1$ or $x \geq 1$	$-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}, y \neq 0$

### (5) EXPONENTIAL FUNCTION ( $a^x$ ):-

An exponential Function is defined by  $F(x) = a^x$  ( $a > 0, a \neq 1$ ), for all  $x \in \mathbb{R}$

$$D_F = \mathbb{R}$$

$$R_F = \mathbb{R}_+$$

Properties

(1)  $a^{x+y} = a^x \cdot a^y$

(2)  $(a^x)^y = a^{xy}$

(3)  $a^x = 1, \Leftrightarrow x = 0$

(4) If  $a > 1, a^x < a^y$  if  $x < y$

If  $a < 1, a^x > a^y$  if  $x < y$

### (5) LOGARITHMIC FUNCTION

The inverse of  $a^x$  is called logarithmic function.

$f(x) = \log_a x$  ( $\log x$  base  $a$ ) is called the logarithmic Function.

$$\text{Dom } f = \mathbb{R}^+, R_f = \mathbb{R}$$

### PROPERTIES

(1)  $\log_a(xy) = \log_a x + \log_a y$

(2)  $\log_a\left(\frac{x}{y}\right) = \log_a x - \log_a y$

(3)  $\log_a x = 0 \Leftrightarrow x = 1$

(4)  $\log_a a = 1$

(5)  $\log_a x = \frac{1}{\log_x a} \quad (x \neq 1)$

(6)  $\log_a x = \log_b x \cdot \log_a b$

(7)  $\log_a x^n = n \log_a x$

(8)  $\log_a \sqrt[n]{x} = \frac{1}{n} \log_a x$

**(6) ABSOLUTE VALUE FUNCTION OR MODULUS**

**FUNCTION ( |x| )** :- The function f defined by  $f(x) = |x| =$   
 $-x$  when  $x < 0$   
 $x$  when  $x \geq 0$

Is called Absolute Value Function.

$D_f = \mathbb{R}$

$R_f = \mathbb{R}^+ \cup \{0\}$

E.g.  $|5| = 5$  (as  $5 > 0$ )

$|-2| = -(-2) = 2$  (as  $-2 < 0$ )

$|0| = 0$

$|3.7| = 3.7$

$|-5.2| = 5.2$

**(8) GREATEST INTEGER FUNCTION OR BRACKET x ( [x] ) :-**

The greatest integer Function [x] is defined as

$$[x] = \begin{cases} x & \text{if } x \in \mathbb{Z} \\ n & \text{if } x \notin \mathbb{Z} \text{ and } n < x < n + 1 \text{ where } n \in \mathbb{Z} \end{cases}$$

Example

$[2] = 2$  (as  $2 \in \mathbb{Z}$ )

$[0] = 0$  (as  $0 \in \mathbb{Z}$ )

$[2.5] = 2$  {as  $2.5 \notin \mathbb{Z}$  and  $2.5$  lies between 2 to 3 i. e.  $2 < 2.5 < 3$ }

$[-1.5] = -2$  {as  $-2 < -1.5 < -1$ }

$[\sqrt{3}] = 1$  { as  $1 < \sqrt{3} < 2$  }

$[-e] = -3$  {as  $2 < e < 3 \Rightarrow -3 < -e < -2$  }

$\text{Dom}[x] = \mathbb{R}$  and  $\text{Range}[x] = \mathbb{Z}$

Functions are categories under two types as follows

**(1) ALGEBRAIC FUNCTION (Three types)**

i) **Polynomial P(x)** =  $a_0 + a_1x + \dots + a_nx^n$

E.g  $F(x) = x^2 + 2x + 3$ ,  $F(x) = 3x + 5$  e

(ii) **Rational Function**  $\left(\frac{P(x)}{Q(x)}\right) = \frac{a_0 + a_1x + \dots + a_nx^n}{b_0 + b_1x + \dots + b_mx^m}$

E.g.  $\frac{x}{x^2+1}$  ,  $\frac{x^2+2x+5}{3x+1}$  are rational functions.

iii) **Irrational Function**  $\{P(x)\}^{p/q}$  e.g.  $\sqrt{x}, (x^2+2x+1)^{2/3}$  etc.

**2) TRANSCENDENTAL FUNCTION**

Trigonometric, logarithmic, exponential functions are called transcendental functions.

Again there are following types of functions as follows.

**EXPLICIT FUNCTION**

$y=f(x)$  i.e. if  $y$  is expressed directly in terms of independent variable  $x$ , then it is called explicit function .

Example  $y =$

$x^2$   $y =$

$2x + 1$  etc.

**IMPLICIT FUNCTION**

Function in which  $x$  and  $y$  cannot be separated from each other (i.e.)  $F(x,y) = 0$  is called implicit Function. E.g.  $x^2+y^2 = 1$   $x^3+2xy+3x^2y^2 = 7$   
are examples implicit functions.

**EVEN FUNCTION:-**

If  $f(-x) = f(x)$ , then  $f(x)$  is called even function.

Example  $f(x) = \cos X$

$f(-x) = \cos (-x) = \cos x$

$=f(x)$  Hence  $f(x) = \cos X$  is an

even function Similarly  $f(x) = x^2, x^4, \dots$

$\dots$  are even functions.

**ODD FUNCTION:-**

If  $f(-x) = -f(x)$ , then  $f(x)$  is called odd function.

$f(x) = \sin x, x, x^3, \dots$  are example of odd functions

**INTRODUCTION TO LIMIT:-**

The concept of limits plays an important role in calculus. Before defining the limit of a function near a point let us consider the following example

Let  $F(x) = \frac{x^2-1}{x-1}$

Now  $F(1) = \frac{1^2-1}{1-1} = \frac{0}{0}$  undefined

But if we take  $x$  close to 1, we obtain different values for  $F(x)$  as follows

TABLE -1

X	0.91	0.93	0.99	0.9999	0.99999
F(X)	1.91	1.93	1.99	1.9999	1.99999

TABLE – 2

X	1.1	1.01	1.001	1.00001	1.000001
F(X)	2.1	2.01	2.001	2.00001	2.000001

In above we can see that when  $x$  gets closer to 1,  $F(x)$  gets closer to 2. However, in this case  $F(x)$  is not defined at  $x=1$ , but as  $x$  approaches to 1  $F(x)$  approaches to 2.

This generates a new concept in setting the value of a function by approach method.

The above value is called limiting value of a Function.

### **SOME DEFINITIONS ASSOCIATED WITH LIMIT:-**

#### **NEIGHBOURHOOD:-**

For every  $a \in \mathbb{R}$ , the open interval  $(a-\delta, a+\delta)$  is called a neighborhood of  $a$  where  $\delta > 0$  is a very very small quantity.

Example  $(1.9, 2.1)$  is a neighborhood of 2. ( $\delta = 0.1$ )

#### **DELETED NEIGHBOURHOOD of 'a':-**

$(a-\delta, a+\delta) - \{a\}$  is called deleted neighborhood of  $a$ .

Left neighborhood of  $a$  is given by  $(a-\delta, a)$ .

Right neighborhood of  $a$  is given by  $(a, a+\delta)$ .

Example

$(1.9, 2.1) - \{2\}$  is a deleted neighborhood of 2.

$(1.9, 2)$  is left neighborhood of 2.

$(2, 2.1)$  is a right neighborhood of 2.

### **DEFINITION OF LIMIT:-**

Given  $\epsilon > 0$ , there exist  $\delta > 0$  depending upon  $\epsilon$  only such that,  $|x-a| < \delta \Rightarrow |f(x)-l| < \epsilon$

$$\text{Then } \lim_{x \rightarrow a} f(x) = l$$

#### EXPLANATION

If for every  $\epsilon > 0$ , we can able to find  $\delta$ , which depends upon  $\epsilon$  only such that  $x \in (a-\delta, a+\delta)$ ,  $\Rightarrow f(x) \in (l-\epsilon, l+\epsilon)$ . In other words when  $x$  gets closer to  $a$  then  $f(x)$  gets closer to  $l$ .

We read  $x \rightarrow a$  as  $x$  tends to 'a' i.e.  $x$  is nearer to  $a$  but  $x \neq a$

$\lim_{x \rightarrow a} f(x)$  limit  $x$  tends a  $f(x)$ . 'l' is called limiting value of  $f(x)$  at  $x = a$ .

In 1<sup>st</sup> example  $\lim_{x \rightarrow 1} \frac{x^2-1}{x-1} = 2$

i.e limiting value of  $f(x)$  at  $x=1$  is 2.

Note:-

Functional value always gives the exact value of a function at a point where as limiting value gives an approximated value of function.

Functional value is either defined or undefined. Similarly limiting value is either exist or does not exist.

### EXISTENCY OF LIMITING VALUE:-

In our first example if we observe table -1 then we see we approach 2 from left in that table.

In table -2 we approach 2 from right.

So in table -1  $x \in (2-\delta, 2)$

And in table -2  $x \in (2, 2+\delta)$

These two approaches give rise to two definitions.

### LEFT HAND LIMIT

When  $x$  approaches  $a$  from left then the value to which  $f(x)$  approaches is called left hand limit of

$f(x)$  at  $x=a$  written as L.H.L. =  $\lim_{x \rightarrow a^-} f(x)$

$x \in a^-$  means  $x \in (a - \delta, a)$ .

### RIGHT HAND LIMIT: -

When  $x$  approaches  $a$  from right then the value to which  $f(x)$  approaches is called right hand limit.

Mathematically

$$\text{R.H.L.} = \lim_{x \rightarrow a^+} f(x)$$

$x \in a^+$  means  $x \in (a, a+\delta)$

### EXISTENCY OF LIMIT

If L.H.L = R.H.L i.e.

Then the limit of  $\lim_{x \rightarrow a} f(x) = l$

$$\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = l$$

the function exists and

Otherwise limit does not exist.

### ALGEBRA OF LIMIT: -

If  $\lim_{x \rightarrow a} f(x) = l$  and  $\lim_{x \rightarrow a} g(x) = m$



Then

$$\begin{aligned} \text{i) } \lim_{x \rightarrow a} \{ f(x) + g(x) \} &= \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x) = l + m \\ \text{ii) } \lim_{x \rightarrow a} \{ f(x) - g(x) \} &= \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x) = l - m \\ \text{iii) } \lim_{x \rightarrow a} \{ f(x) \cdot g(x) \} &= \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x) = l \cdot m \\ \text{iv) } \lim_{x \rightarrow a} \left\{ \frac{f(x)}{g(x)} \right\} &= \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)} = \frac{l}{m} \quad (\text{provided } m \neq 0) \\ \text{v) } \lim_{x \rightarrow a} K &= K \quad (K \text{ is constant}) \\ \text{vi) } \lim_{x \rightarrow a} k f(x) &= k \lim_{x \rightarrow a} f(x) = k l \\ \text{(vii) } \lim_{x \rightarrow a} \log_b f(x) &= \log_b \lim_{x \rightarrow a} f(x) = \log_b l \\ \text{(viii) } \lim_{x \rightarrow a} e^{f(x)} &= e^{\lim_{x \rightarrow a} f(x)} = e^l \\ \text{(ix) } \lim_{x \rightarrow a} f(x)^n &= \{ \lim_{x \rightarrow a} f(x) \}^n = l^n \\ \text{(x) } \lim_{x \rightarrow a} f(x) &= \lim_{y \rightarrow a} f(y) = l \\ \text{(xi) } \lim_{x \rightarrow a} |f(x)| &= \left| \lim_{x \rightarrow a} f(x) \right| = |l| \\ \text{(xii) If } \lim_{x \rightarrow a} f(x) &= \infty, \text{ then } \lim_{x \rightarrow a} \frac{1}{f(x)} = 0 \end{aligned}$$

### EVALUATION OF LIMIT:-

When we evaluate limits it is not necessary to test the existency of limit always. So in this section we will discuss various methods of evaluating limits.

#### (1) EVALUATION OF ALGEBRAIC LIMITS :-

(2) Method -> (i) Direct substitution (ii) Factorisation (iii) Rationalisation

#### i) Direct Substitution :-

If  $f(x)$  is an algebraic function and  $f(a)$  is finite. Then  $\lim_{x \rightarrow a} f(x)$  is equal to  $f(a)$

i.e. we can substitute  $x$  by  $a$ .

Let us consider following examples.

#### Example -1

Evaluate  $\lim_{x \rightarrow 0} (x^2 + 2x + 1)$

ANS  $\square$

$$\lim_{x \rightarrow 0} (x^2 + 2x + 1) = 0^2 + 2 \cdot 0 + 1 = 1$$

#### Example -2

Evaluate  $\lim_{x \rightarrow -1} \frac{x-1}{x^2+2x-1}$

ANS  $\square$

$$\lim_{x \rightarrow -1} \frac{x-1}{x^2+2x-1} = \frac{(-1)-1}{(-1)^2+2 \times (-1)-1} = \frac{-2}{1-2-1} = \frac{-2}{-2} = 1$$

**Example – 3**

$$\lim_{x \rightarrow 1} \frac{\sqrt{x}}{\sqrt{x+2}} = ?$$

Ans :-

$$\lim_{x \rightarrow 1} \frac{\sqrt{x}}{\sqrt{x+2}} = \frac{\sqrt{1}}{\sqrt{1+2}} = \frac{1}{\sqrt{3}}$$

**Example -4**

Evaluate  $\lim_{x \rightarrow 1} \frac{x^2-1}{x-1} = \frac{1^2-1}{1-1} = \frac{0}{0}$ , Which cannot be determined.

**NOTE: -**

So here direct substitution method fails to find the limiting value. In this case we apply following method.

**ii) FACTORISATION METHOD :-**

If the given Function is a rational function  $\frac{f(x)}{g(x)}$ , and  $\frac{f(a)}{g(a)}$  is in  $\frac{0}{0}$  form then we apply factorisation method i.e we factorise f(x) and g (x) and cancel the common factor. After cancellation we again apply direct substitution, if result is a finite number otherwise we repeat the process .

This method is clearly explained in following example.

**Example -4**

Evaluate  $\lim_{x \rightarrow 1} \frac{x^2-1}{x-1}$

Ans :-  $\lim_{x \rightarrow 1} \frac{x^2-1}{x-1} = \lim_{x \rightarrow 1} \frac{(x+1)(x-1)}{(x-1)}$

$$= \lim_{x \rightarrow 1} (x+1) \quad \{x \neq 1 \text{ means } x \neq 1 \Rightarrow (x-1) \neq 0\}$$

$$= 1+1 = 2 \quad \{\text{after cancellation we can apply the direct substitution}\}$$

**Example -5**

Evaluate  $\lim_{x \rightarrow -3} \frac{x^2+7x+12}{x^2+5x+6}$

ANS :-

$$\lim_{x \rightarrow -3} \frac{x^2+7x+12}{x^2+5x+6} \quad \{\text{by putting } x=-3 \text{ we can easily check that the question is in } \frac{0}{0} \text{ form}\}$$

$$= \lim_{x \rightarrow -3} \frac{x^2+4x+3x+12}{x^2+2x+3x+6}$$

$$= \lim_{x \rightarrow -3} \frac{x(x+4)+3(x+4)}{x(x+2)+3(x+2)}$$

$$= \lim_{x \rightarrow -3} \frac{(x+4)(x+3)}{(x+2)(x+3)} \quad \{x \neq -3 \text{ then } x+3 \neq 0\}$$

$$= \lim_{x \rightarrow -3} \frac{(x+4)}{(x+2)} = \frac{-3+4}{-3+2} = \frac{1}{-1} = -1$$

**iii) Rationalisation method :-**

When either the numerator or the denominator contain some irrational functions and direct substitution gives  $\frac{0}{0}$  form, then we apply rationalisation method. In this method we rationalize the irrational function to eliminate the  $\frac{0}{0}$  form. This can be better explained in following examples.

**Example -7**

Evaluate  $\lim_{x \rightarrow 0} \frac{x}{\sqrt{x+1}-1}$

ANS :-

$\lim_{x \rightarrow 0} \frac{x}{\sqrt{x+1}-1}$  { In order to rationalize  $\sqrt{x+1}-1$  we have to apply  $a^2-b^2$  formula  
 $a^2-b^2 = (a+b)(a-b)$  so here  $a-b$  is present, so we have to

multiply  $a+b$  i.e.  $\sqrt{x+1}+1$  }

$$\begin{aligned} &= \lim_{x \rightarrow 0} \frac{x(\sqrt{x+1}+1)}{(\sqrt{x+1}-1)(\sqrt{x+1}+1)} \\ &= \lim_{x \rightarrow 0} \frac{x(\sqrt{x+1}+1)}{\sqrt{x+1}^2-1^2} \\ &= \lim_{x \rightarrow 0} \frac{x(\sqrt{x+1}+1)}{x+1-1} = \lim_{x \rightarrow 0} \frac{x(\sqrt{x+1}+1)}{x} \\ &= \lim_{x \rightarrow 0} (\sqrt{x+1}+1) = \sqrt{0+1}+1 = 1+1 = 2 \end{aligned}$$

**(3) Evaluating limit when  $x \rightarrow \infty$**

In order to evaluate infinite limits we use some formulas and techniques.

Formulas (i)  $\lim_{x \rightarrow \infty} x^n = \infty, n > 0$

(ii)  $\lim_{x \rightarrow \infty} \frac{1}{x^n} = 0, n > 0$

$\frac{f(x)}{g(x)}$

When we evaluate functions in  $\frac{f(x)}{g(x)}$  form, then we use the following technique

Divide both  $f(x)$  and  $g(x)$  by  $x^k$  where  $x^k$  is the highest order term in  $g(x)$ .

It can be better understood by following examples.

**Example - 9**

Evaluate  $\lim_{x \rightarrow \infty} \frac{3x^2+x-1}{2x^2-7x+5}$

ANS:-  $\lim_{x \rightarrow \infty} \frac{3x^2+x-1}{2x^2-7x+5}$

{Dividing numerator and denominator by highest order term in denominator i.e.  $x^2$ }

$$\begin{aligned} &= \lim_{x \rightarrow \infty} \frac{\frac{3x^2+x-1}{x^2}}{\frac{2x^2-7x+5}{x^2}} \\ &= \lim_{x \rightarrow \infty} \frac{\frac{3x^2}{x^2} + \frac{x}{x^2} - \frac{1}{x^2}}{\frac{2x^2}{x^2} - \frac{7x}{x^2} + \frac{5}{x^2}} = \lim_{x \rightarrow \infty} \frac{3 + \frac{1}{x} - \frac{1}{x^2}}{2 - \frac{7}{x} + \frac{5}{x^2}} \\ &\qquad\qquad\qquad \frac{1}{1} \qquad\qquad\qquad \frac{1}{1} \end{aligned}$$

$$= \frac{\lim_{x \rightarrow \infty} 3 + \lim_{x \rightarrow \infty} x - \lim_{x \rightarrow \infty} x^2}{\lim_{x \rightarrow \infty} 2 - \lim_{x \rightarrow \infty} \frac{7}{x} + \lim_{x \rightarrow \infty} \frac{5}{x^2}} \text{ (applying algebra of limits)}$$

$$= \frac{3+0-0}{2-0+0} = \frac{3}{2}$$

**Important note in  $\infty$  limit evaluation:-**

$$\lim_{x \rightarrow \infty} \frac{a_0 + a_1x + \dots + a_mx^m}{b_0 + b_1x + b_2x^2 + \dots + b_nx^n} = \begin{cases} \frac{m}{b_n} & \text{if } m = n \\ 0 & \text{if } m < n \\ \infty & \text{if } m > n \end{cases}$$

**Example-13**

If  $\lim_{x \rightarrow \infty} \left( \frac{x^2-1}{x+1} - ax - b \right) = 2$ , find the values of  $a$  and  $b$ .

Solution -> Given  $\lim_{x \rightarrow \infty} \left( \frac{x^2-1}{x+1} - ax - b \right) = 2$

$$\Rightarrow \lim_{x \rightarrow \infty} \left( \frac{x^2-1-ax^2-ax-bx-b}{x+1} \right) = 2$$

$$\Rightarrow \lim_{x \rightarrow \infty} \frac{x^2(1-a) - x(a+b) - (b+1)}{x+1} = 2$$

As result is finite non zero quantity

$\Rightarrow$  Degree of numerator polynomial = degree of denominator polynomial

$$\Rightarrow \boxed{\text{Degree of polynomial in numerator} = 1}$$

{As  $x+1$  has degree = 1}

$$\Rightarrow 1-a = 0 \Rightarrow \boxed{a=1}$$

Now putting  $a = 1$  in above evaluation

$$\lim_{x \rightarrow \infty} \frac{-x(1+b) - (b+1)}{x+1} = 2$$

$$\Rightarrow \frac{-(1+b)}{1} = 2 \quad \{\text{by important note}\}$$

$$\Rightarrow \boxed{-1-b=2} \quad \left\{ \lim_{x \rightarrow \infty} \frac{a_0 + a_1x + \dots + a_mx^m}{b_0 + b_1x + \dots + b_nx^n} = \frac{a_m}{b_n} \text{ where } m = n \right\}$$

$$\Rightarrow \boxed{b=-1-2=-3}$$

Therefore  $a=1$  and  $b=-3$

**(4) Important Formulas in limit**

(1)  $\lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = na^{n-1}$  where  $a > 0$  and  $n \in \mathbb{R}$

(2)  $\lim_{x \rightarrow 0} \frac{a^x - 1}{x} = \log_{e a}$

In particular  $\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1$

$$(3) \lim_{x \rightarrow 0} (1+x)^{\frac{1}{x}} = e$$

$$(4) \lim_{x \rightarrow \infty} (1 + \frac{1}{x})^x = e$$

$$(5) \lim_{x \rightarrow 0} \frac{\log_a(1+x)}{x} = \log_a e$$

In particular  $\lim_{x \rightarrow 0} \frac{\log_e(1+x)}{x} = \log_e e = 1$

$$(6) \lim_{x \rightarrow 0} \cos x = 1$$

$$(7) \lim_{x \rightarrow 0} \sin x = 0$$

$$(8) \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$$

$$(9) \lim_{x \rightarrow 0} \frac{\tan x}{x} = 1$$

**SUBSTITUTION METHOD:-**

In order to apply known formula sometimes we apply substitution method. In this method x is replaced by another variable u, and then we apply formula on ‘u’.

Let us consider the following example.

**Example – 14:-**

Evaluate  $\lim_{x \rightarrow 0} \frac{\sin 2x}{x}$

ANS:- Let  $2x=u \Rightarrow$  when  $x \rightarrow 0$

$$u \rightarrow 0 \text{ (as } u = 2x)$$

$$\begin{aligned} \text{Now } \lim_{x \rightarrow 0} \frac{\sin 2x}{x} &= \lim_{u \rightarrow 0} \frac{\sin u}{\frac{u}{2}} = 2 \lim_{u \rightarrow 0} \frac{\sin u}{u} \\ &= 2 \times 1 = 2 \end{aligned}$$

In general

Putting  $\lambda x = u$

$\lim_{x \rightarrow 0} \frac{f(\lambda x)}{x} = \lim_{u \rightarrow 0} \frac{f(u)}{\frac{u}{\lambda}}$
---

Hence some of the formulas may be stated as follows

$$1) \lim_{x \rightarrow 0} \frac{a^{\lambda x} - 1}{\lambda x} = \log_a e$$

In particular  $\lim_{x \rightarrow 0} \frac{e^{\lambda x} - 1}{\lambda x} = 1$

$$2) \lim_{x \rightarrow 0} (1 + \lambda x)^{\frac{1}{\lambda x}} = e$$

$$3) \lim_{x \rightarrow \infty} (1 + \frac{1}{\lambda x})^{\lambda x} = e$$

$$4) \lim_{x \rightarrow 0} \frac{\log_a(1+\lambda x)}{\lambda x} = \log_a e$$

In particular  $\lim_{x \rightarrow 0} \frac{\log_e(1+\lambda x)}{\lambda x} = 1$

$$5) \lim_{x \rightarrow 0} \frac{\sin \lambda x}{\lambda x} = 1$$

$$6) \lim_{x \rightarrow 0} \frac{\tan \lambda x}{\lambda x} = 1$$

**Some examples based on the formulas**

(1) Evaluate  $\lim_{x \rightarrow 0} \frac{\sin 3x}{\tan 5x}$

Ans :-

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{\sin 3x}{\tan 5x} &= \lim_{x \rightarrow 0} \frac{\frac{3 \sin 3x}{3x}}{\frac{5 \tan 5x}{5x}} \\ &= \frac{3}{5} \lim_{x \rightarrow 0} \frac{\left(\frac{\sin 3x}{3x}\right)}{\left(\frac{\tan 5x}{5x}\right)} = \frac{3}{5} \times \frac{1}{1} = \frac{3}{5} \end{aligned}$$

(2) Evaluate  $\lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2}$  (2014 S)

$$\begin{aligned} \text{Ans :- } \lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} &= \lim_{x \rightarrow 0} \frac{2 \sin^2 \frac{x}{2}}{x^2} \\ &= \lim_{x \rightarrow 0} 2 \cdot \frac{\sin \frac{x}{2}}{\frac{x}{2}} \cdot \frac{\sin \frac{x}{2}}{\frac{x}{2}} = \frac{1}{2} \lim_{x \rightarrow 0} \frac{\sin \frac{x}{2}}{\frac{x}{2}} \cdot \frac{\sin \frac{x}{2}}{\frac{x}{2}} = \frac{1}{2} \times 1 \times 1 = \frac{1}{2} \end{aligned}$$

**Exercise**

**1. Evaluate the following limits(2 marks)**

(i)  $\lim_{x \rightarrow 0} \frac{\sqrt{x+1} - 1}{x}$

(ii)  $\lim_{x \rightarrow 0} \frac{\sin ax}{\sin bx}$  (a;b ≠ 0)

(iii)  $\lim_{x \rightarrow 0} \frac{\sin 3x}{\sin 2x}$

**2. Evaluate the following limits (5 marks)**

(i)  $\lim_{x \rightarrow 1} \frac{2^{x-1} - 1}{\sqrt{x} - 1}$

(ii)  $\lim_{x \rightarrow 5} \frac{\sqrt{x-1} - 2}{x-5}$

(iii)  $\lim_{x \rightarrow 1} \frac{\frac{1}{x^m} - 1}{\frac{1}{x^n} - 1}$

(iv)  $\lim_{n \rightarrow \infty} \frac{1^3 + 2^3 + 3^3 + \dots + n^3}{n^4}$

(v)  $\lim_{x \rightarrow \infty} x^2 \{ \sqrt{x^4 + a^2} - \sqrt{x^4 - a^2} \}$

(xii)  $\lim_{x \rightarrow 0} \frac{\sqrt[3]{1+x} - \sqrt[3]{1-x}}{x}$

$$\begin{aligned}
 & \text{(xiii)} \quad \lim_{x \rightarrow 0} \frac{a^x - b^x}{c^x - d^x} \\
 \text{(2016-S)} \quad & \text{(xiv)} \quad \lim_{x \rightarrow 0} \frac{\cos 2x - \cos 3x}{x^2} \\
 & \text{(2017-S)} \\
 \text{(xv)} \quad & \lim_{x \rightarrow 0} \frac{\sqrt{3-2x} - \sqrt{3}}{x} \\
 \text{(xvi)} \quad & \lim_{x \rightarrow 0} \frac{\sqrt{1+x} - \sqrt{1-x}}{\sin^{-1} x} \qquad \qquad \qquad \text{(2017-w)}
 \end{aligned}$$

**Continuity and Discontinuity of Function** In the figure we observe that the 1<sup>st</sup> graph of a function in Fig-1 can be drawn on a paper without raising pencil i.e. 1<sup>st</sup> graph is continuously moving where as Fig -2 represents a graph , which cannot be drawn without raising the pencil. Because there are gaps or breaks. So, it is discontinuous.




Fig-1




Fig-2

The feature of the graph of a function displays an important property of the function called continuity of a function.

**Continuity of a Function at a point**

Definition – A function f(x) is said to be continuous at x = a, if it satisfies the following conditions

- (i)  $\lim_{x \rightarrow a} f(x)$  exists.
- (ii) f(a) is defined i.e. finite
- (iii)  $\lim_{x \rightarrow a} f(x) = f(a)$

If one or more of the above condition fail, the function f(x) is said to be discontinuous at x= a.

**Continuous Function**

A function is said to be continuous if it is continuous at each point of its domain. **Working**

**procedure for testing continuity at a point x = a**

**1<sup>st</sup> step** – First find  $\lim_{x \rightarrow a} f(x)$  by using concepts from previous chapter.

If  $\lim_{x \rightarrow a} f(x)$  does not exist then, f(x) is discontinuous at x = a.

If  $\lim_{x \rightarrow a} f(x) = l$ , then go to 2<sup>nd</sup> step.

**2<sup>nd</sup> step** – Find f(a) from the given data

If  $f(a)$  is undefined then  $f(x)$  is not continuous at  $x = a$ .

If  $f(a)$  has finite value then go to 3<sup>rd</sup> step.

**3<sup>rd</sup> step** – Compare  $\lim_{x \rightarrow a} f(x)$  and  $f(a)$

If  $\lim_{x \rightarrow a} f(x) = f(a)$ , then  $f(x)$  is continuous at  $x = a$ , otherwise  $f(x)$  is discontinuous at  $x = a$ .

**Examples**

**Q1.** Examine the continuity of the function  $f(x)$  at

$$x = 3. \quad f(x) = \begin{cases} \frac{x^2-9}{x-3} & x \neq 3 \\ 6 & x = 3 \end{cases}$$

Ans:-

$$\lim_{x \rightarrow 3} f(x) = \lim_{x \rightarrow 3} \frac{x^2-9}{x-3} = \lim_{x \rightarrow 3} \frac{(x+3)(x-3)}{(x-3)} = \lim_{x \rightarrow 3} (x+3) = 3+3 = 6 \quad \{ \text{As } x \neq 3, x \neq 3 \Rightarrow x-3 \neq 0 \}$$

From given data  $f(3) = 6$

Now from above  $\lim_{x \rightarrow 3} f(x) = f(3)$

Therefore,  $f(x)$  is continuous at  $x = 3$ .

**Q2.** Test continuity of  $f(x)$  at ‘0’ where,

$$f(x) = \begin{cases} (1+3x)^{\frac{1}{3x}} & x \neq 0 \\ e^3 & x = 0 \end{cases}$$

Ans:-  $\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} (1+3x)^{\frac{1}{3x}}$

$$= \lim_{x \rightarrow 0} (1+3x)^{\frac{1}{3x} \cdot 3}$$

$$= \lim_{x \rightarrow 0} \{ (1+3x)^{\frac{1}{3x}} \}^3$$

$$= \{ \lim_{x \rightarrow 0} (1+3x)^{\frac{1}{3x}} \}^3$$

$$= e^3$$

{ As  $\lim_{x \rightarrow 0} (1+x)^{\frac{1}{x}} = e$

$$\lim_{x \rightarrow 0} (1+\lambda x)^{\frac{1}{\lambda x}} = e$$

In particular  $\lim_{x \rightarrow 0} (1+3x)^{\frac{1}{3x}} = e$  and we know,  $\lim_{x \rightarrow a} \{f(x)\}^n = \{ \lim_{x \rightarrow a} f(x) \}^n$

From given data  $f(0) = e^3$

Hence,  $\lim_{x \rightarrow 0} f(x) = f(0)$



Therefore,  $f(x)$  is continuous at  $x = 0$ .

**Q3.** Test continuity of  $f(x)$  at  $x = 0$

$$f(x) = \begin{cases} \frac{|x|}{x} & x \neq 0 \\ 0 & x = 0 \end{cases}$$

Ans.  $\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \frac{|x|}{x}$

As  $|x|$  is present and  $x \neq 0$ , so we have to evaluate the above limit by L.H.L and R.H.L

method L.H.L =  $\lim_{x \rightarrow 0^-} \frac{|x|}{x} \quad \{ x \rightarrow 0^- \Rightarrow x < 0 \}$

$$= \lim_{x \rightarrow 0^-} \frac{-x}{x}$$

$$= \lim_{x \rightarrow 0^-} (-1) = (-1)$$

R.H.L =  $\lim_{x \rightarrow 0^+} \frac{|x|}{x} \quad \{ x \rightarrow 0^+ \Rightarrow x > 0 \}$

$$= \lim_{x \rightarrow 0^+} \frac{x}{x} = \lim_{x \rightarrow 0^+} (1) = 1$$

Hence, L.H.L  $\neq$  R.H.L

Therefore,  $f(x)$  does not exist.

Hence  $f(x)$  is not continuous at  $x = 0$ .

## UNIT-III

# DERIVATIVES

## Introduction

---

The study of differential calculus originated in the process of solving the following three problems

1. From the astronomical consideration particularly involving an attempt to have a better approximation of  $\pi$  as developed by Bhaskaracharya, Madhava and Nilakantha.
2. Finding the tangent to any arbitrary curve as developed by Fermat and Leibnitz.
3. Finding rate of change as developed by Fermat and Newton.

In this chapter we define derivative of a function, give its geometrical and physical interpretation and discuss various laws of derivatives etc.

## Objectives

---

After studying this lesson, you will be able to:

- (1) Define and Interpret geometrically the derivative of a function  $y = f(x)$  at  $x = a$ .
- (2) State derivative of some standard function.
- (3) Find the derivative of different functions like composite function, implicit function using different techniques.
- (4) Find higher order derivatives of a particular function by successive differentiation method.
- (5) Determine rate of change and tangent to a curve.
- (6) Find partial derivative of a function with more than one variable with respect to variables.
- (7) Define Euler's theorem and apply it solve different problems based on partial differentiation.

## Expected background knowledge

---

1. Function
2. Limit and continuity of a function at a point.

## Derivative of a function

---

Consider a function  $y = x^2$

**Table-1**

x	5	5.1	5.01	5.001	5.0001
y	25	26.01	25.1001	25.010001	25.00100001

Let  $x = 5$  and  $y = 25$  be a reference point

We denote the small changes in the value of  $x$  as ' $\delta x$ ',

$\delta x$  = small change in x  $\delta y$  = change in y,

when there is a change of  $\delta x$  in x.

Now,  $\frac{\delta y}{\delta x}$  is called Increment ratio or Newton quotient or average rate of change of y.

Now, let us write table -1 in terms of  $\delta x$ ,  $\delta y$  as

**Table-2**

$\delta x$	0.1	0.01	0.001	0.0001
$\delta y$	1.01	0.1001	0.010001	0.00100001
$\frac{\delta y}{\delta x}$	10.1	10.01	10.001	10.0001

From table-2  $\delta y$  varies as  $\delta x$  varies

It is clear from the table when  $\delta x \rightarrow 0$  and  $\frac{\delta y}{\delta x} \rightarrow 10$

This  $\frac{\delta y}{\delta x}$  when  $\delta x \rightarrow 0$  is the instantaneous rate of change of y at the value of x.

In above case  $x = 5$ , so  $\frac{dy}{dx}$  at  $x = 5$  is 10

**Definition of derivative of a function (Differentiation)**

If  $y = f(x)$  is a function. Then derivative of y with respect to x is given by

$$\frac{dy}{dx} = \frac{\delta y}{\delta x}$$

$\frac{dy}{dx}$  is also denoted by  $f'(x)$

$$\frac{dy}{dx} = \frac{\delta y}{\delta x} = f'(x) = f'$$

are same notations

**Process of finding derivatives of dependent variable w.r.t. independent variable is called differentiation.**

**Derivative of a function at a point 'a'**

Derivative of  $y = f(x)$  at a point 'a' in the domain  $D_f$  is given by

$$\left. \frac{dy}{dx} \right|_{x=a} = f'(a) = \underline{\hspace{2cm}}$$


**Example -1**

Find the derivative of  $f(x) = x^2$  at  $x = 5$

$$\begin{aligned} \text{Ans. } \left. \frac{dy}{dx} \right|_{x=5} = f'(5) &= \lim_{h \rightarrow 0} \frac{f(5+h) - f(5)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(5+h)^2 - 5^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{(5+h+5)(5+h-5)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(10+h)h}{h} = \lim_{h \rightarrow 0} (10 + h) = 10 \end{aligned}$$

**Geometrical Interpretation of  $\frac{dy}{dx}$**

(Fig.-1)



Let  $f(x)$  is represented by the curve in fig-1 given above.

Let  $Q(x+\delta x, y + \delta y)$  be the neighbourhood of  $P(x,y)$ .  $PM$  and  $QN$  are drawn perpendicular to Xaxis.

$PS \perp QN$

Let QP Secant meets x-axis, (by extending it) and  $\overline{QP}$  make angle  $\theta$  with x-axis then angle QPS =  $\theta$

In  $\Delta QPS$ ,  $\tan \theta = \frac{QS}{PS} = \frac{\delta y}{\delta x}$

As  $QN = y + \delta y$ ,  $NS = PM = y$

$\Rightarrow QS = QN - NS = \delta y$ .

Similarly,  $ON = x + \delta x$  and  $OM = x \Rightarrow PS = MN = ON - OM$

=  $\delta x$  When  $\delta x \rightarrow 0$  then  $Q \rightarrow P$  and QP secant becomes tangent at

In PQS  $\theta = \frac{\delta y}{\delta x}$  { gives slope of PQ line }  $\Delta$

We know

$$\frac{\delta y}{\delta x} = \tan \theta$$

P.

Now when  $\delta x \rightarrow 0$  the line PQ becomes tangent at P

So,

$$\frac{dy}{dx} = \tan \theta = \text{slope of the tangent to the curve at P.}$$

So derivative of a function at a point represents the slope or gradient of the tangent at that point.

**Example 2**

Q. Find the slope of the tangent to the curve  $y = x^2$  at  $x = 5$ .

**Ans.** As we have done it in example – 1.

$$\left. \frac{dy}{dx} \right|_{x=5} = 10$$

Therefore, slope of the tangent at  $x = 5$  is 10.

**Derivative of some standard functions**

---

1.  $\frac{d}{dx}(c) = 0$

2.  $\frac{d}{dx}(x^n) = n \cdot x^{n-1}$
3.  $\frac{d}{dx}(a^x) = a^x \log_e a$       In particular  $\frac{d}{dx}(e^x) = e^x$  4.      In particular  $\frac{d}{dx}(\log_e x) = \frac{1}{x}$   
 $\frac{d}{dx}(\log_a x) = \frac{1}{x \log_e a}$        $(\ln x) = \frac{1}{x}$
5.  $\frac{d}{dx}(\sin x) = \cos x$
6.  $\frac{d}{dx}(\cos x) = -\sin x$
7.  $\frac{d}{dx}(\tan x) = \sec^2 x$
8.  $\frac{d}{dx}(\cot x) = -\operatorname{cosec}^2 x$
9.  $\frac{d}{dx}(\sec x) = \sec x \cdot \tan x$
10.  $\frac{d}{dx}(\operatorname{cosec} x) = -\operatorname{cosec} x \cdot \cot x$
11.  $\frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}$
12.  $\frac{d}{dx}(\cos^{-1} x) = -\frac{1}{\sqrt{1-x^2}}$
13.  $\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}$
14.  $\frac{d}{dx}(\cot^{-1} x) = -\frac{1}{1+x^2}$
15.  $\frac{d}{dx}(\sec^{-1} x) = \frac{1}{|x|\sqrt{x^2-1}}$
16.  $\frac{d}{dx}(\operatorname{cosec}^{-1} x) = -\frac{1}{|x|\sqrt{x^2-1}}$

### Algebra of derivatives or fundamental theorems of derivatives

---

If  $f(x)$  and  $g(x)$  are both derivable functions i.e. their derivative exists then,

- (i)  $\frac{d}{dx}\{cf(x)\} = c f'(x)$
- (ii)  $\frac{d}{dx}(f + g) = f' + g'$
- (iii)  $\frac{d}{dx}(f - g) = f' - g'$
- (iv)  $\frac{d}{dx}\{fg\} = fg' + f'g$
- (v)  $\frac{d}{dx}\left\{\frac{f}{g}\right\} = \frac{f'g - fg'}{g^2}$

#### Example-3

Find the derivative of the following:

(i)  $3x^3$

(ii)  $6\sqrt{x}$

(iii)  $9 \cdot 3^x$

(iv)  $5 \cot x$

**Ans.**

(i) 
$$\frac{dy}{dx} = \frac{d}{dx} (3x^3) = 3 \frac{d(x^3)}{dx} = 3 \times 3 x^{3-1} = 9x^2$$

(ii) 
$$\frac{dy}{dx} = \frac{d(6\sqrt{x})}{dx} = 6 \frac{d(\sqrt{x})}{dx} = 6 \frac{d(x^{\frac{1}{2}})}{dx} = 6 \cdot \frac{1}{2} x^{\frac{1}{2}-1} = 6 \cdot \frac{1}{2} x^{-\frac{1}{2}} = \frac{3}{\sqrt{x}}$$

(iii) 
$$\frac{dy}{dx} = \frac{d(9 \cdot 3^x)}{dx} = 9 \frac{d(3^x)}{dx} = 9 \cdot 3^x \ln 3$$

(iv) 
$$\frac{d(5 \cot x)}{dx} = 5 \frac{d(\cot x)}{dx} = 5 (-\operatorname{cosec}^2 x) = -5 \operatorname{cosec}^2 x$$

**Derivative of a composite function (Chain Rule)****Composite function**

A function formed by composition of more than one function is called composite function.

**Example of composite functions**

1)  $\sin x^2$  is form by composition of two functions, one is  $\sin x$  function and other is  $x^2$ .

$$y = \sin x^2 = \sin u \text{ where } u = x^2$$

2) Similarly  $y = \sqrt{x^2 + 3x + 1}$  is written as

$$y = \sqrt{u} \text{ where } u = x^2 + 3x + 1$$

3)  $y = \sqrt{\sin(x^2 + 1)}$  is form by composition of three functions.  $y = \sqrt{u}$  where  $u = \sin v$  and  $v = (x^2 + 1)$

**Chain Rule**

If  $y = f(u)$  and  $u$  is a function of  $x$  defined by  $u = g(x)$ , then

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$$

**Generalized chain rule**

If  $y$  is a differentiable function of  $u$ ,  $u$  is a differentiable function of  $v$ , ..... and finally  $v$  is a differentiable function of  $x$ . Then

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dx} \dots \dots$$

### Example – 9

Find  $\frac{dy}{dx}$

(i).  $y = (x^2 + 2x - 1)^5$

(ii)  $y = \cot^3 x$

Ans.

(i)  $y = (x^2 + 2x - 1)^5$

Here,  $y = u^5$  and  $u = x^2 + 2x - 1$

$$\frac{du}{dx} = 2x + 2 = 2(x+1) \quad \text{and} \quad \frac{dy}{du} = 5u^4$$

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$$

$$= 5u^4 \cdot 2(x+1)$$

$$= 10(x^2 + 2x - 1)^4 (x+1)$$

(ii)  $y = \cot^3 x$  can be written as  $y = u^3$

where  $u = \cot x$

$$\frac{du}{dx} = -\operatorname{cosec}^2 x, \quad \frac{dy}{du} = 3u^2$$

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} = 3u^2(-\operatorname{cosec}^2 x)$$

$$= -3 \cot^2 x \operatorname{cosec}^2 x$$

### Methods of differentiation

---

We use following two methods for differentiation of some functions.

(i) Substitution

(ii) Use of logarithms

### Substitution



Sometimes with proper substitution we can transform the given function to a simpler function in the new variable so that the differentiation w.r.t to new variable becomes easier. After differentiation we again re-substitute the old variable. This can be better understood by following examples.

**Note**

*When we apply substitution method, then we must have proper knowledge about trigonometric formulae. Because it makes the choice of new variable easy. If proper substitution is not made, then problem will be more complicated than original.*

**Example –15**

If  $y = \sec^{-1}\left(\frac{\sqrt{a^2+x^2}}{a}\right)$  then find  $\frac{dy}{dx}$

**Ans.**

$$\begin{aligned} y &= \sec^{-1}\left(\frac{\sqrt{a^2+x^2}}{a}\right) && \text{Put } x = a \tan \theta \\ &= \sec^{-1}\left(\frac{\sqrt{a^2+a^2\tan^2\theta}}{a}\right) && = \sec^{-1}\left(\frac{\sqrt{a^2(1+\tan^2\theta)}}{a}\right) \\ &= \sec^{-1}\left(\frac{\sqrt{a^2\sec^2\theta}}{a}\right) && = \sec^{-1}\left(\frac{a\sec\theta}{a}\right) \\ &= \sec^{-1}(\sec\theta) = \theta && = \tan^{-1}\left(\frac{x}{a}\right) \end{aligned}$$

$$\begin{aligned} \text{Now } \frac{dy}{dx} &= \frac{d}{dx} \left\{ \tan^{-1}\left(\frac{x}{a}\right) \right\} = \frac{1}{1+\left(\frac{x}{a}\right)^2} \frac{d}{dx} \left(\frac{x}{a}\right) \\ &= \frac{1}{\left(1+\frac{x^2}{a^2}\right)} \left(\frac{1}{a}\right) = \frac{1}{a} \frac{1}{\frac{a^2+x^2}{a^2}} \\ &= \frac{a^2}{a(x^2+a^2)} = \frac{a}{x^2+a^2} \end{aligned}$$

**Example – 16**

Differentiate  $\sin^2(\cot^{-1} \sqrt{\frac{1+x}{1-x}})$  w.r.t x. **[2018-**

**S] Ans.**

$$\begin{aligned} y &= \sin^2(\cot^{-1} \sqrt{\frac{1+x}{1-x}}) && \{ \text{Put } x = \cos 2\theta \Rightarrow \theta = \frac{\cos^{-1} x}{2} \} \\ &= \sin^2(\cot^{-1} \sqrt{\frac{1+\cos 2\theta}{1-\cos 2\theta}}) = \sin^2 \left( \cot^{-1} \sqrt{\frac{2\cos^2\theta}{2\sin^2\theta}} \right) \end{aligned}$$

$$= \sin^2 (\cot^{-1} \sqrt{\cot^2 \theta}) = \sin^2 \cot^{-1}(\cot \theta) = \sin^2 \theta$$

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{d\theta} \cdot \frac{d\theta}{dx} = \frac{d}{d\theta} (\sin^2 \theta) \frac{d}{dx} \left( \frac{\cos^{-1} x}{2} \right) \\ &= 2 \sin \theta \cos \theta \cdot \frac{1}{2} \left( \frac{-1}{\sqrt{1-x^2}} \right) \\ &= \sin 2\theta \left( \frac{-1}{2\sqrt{1-x^2}} \right) = -\frac{\sqrt{1-\cos^2 2\theta}}{2\sqrt{1-x^2}} \\ &= -\frac{\sqrt{1-x^2}}{2\sqrt{1-x^2}} = -\frac{1}{2} \text{ (Ans)} \end{aligned}$$

### Differentiation using logarithm

---

When a function appears as an exponent of another function we make use of logarithms.

#### Example – 18

Differentiate  $(\sin x)^{\tan x}$

**Ans.**

$$y = (\sin x)^{\tan x}$$

Taking logarithms of both sides we have,

$$\ln y = \ln(\sin x)^{\tan x}$$

$$\Rightarrow \ln y = \tan x \cdot \ln \sin x$$

Differentiating both sides w.r.t  $x$ , we have

$$\Rightarrow \frac{1}{y} \frac{dy}{dx} = \tan x \cdot \frac{1}{\sin x} \cdot \cos x + \sec^2 x \cdot \ln \sin x$$

$$\Rightarrow \frac{1}{y} \frac{dy}{dx} = \tan x \cdot \cot x + \sec^2 x \cdot \ln \sin x = 1 + \sec^2 x \cdot \ln \sin x$$

$$\Rightarrow \frac{dy}{dx} = y (1 + \sec^2 x \cdot \ln \sin x)$$

$$\text{Hence } \frac{dy}{dx} = (\sin x)^{\tan x} (1 + \sec^2 x \cdot \ln \sin x)$$

#### Example – 19

Differentiate  $y$

$$= \frac{\sqrt{\quad}}{\quad}$$

**Ans.**

$$y = \frac{\sqrt{\quad}}{\quad}$$

Taking logarithm of both sides

$$\Rightarrow \ln y = \ln(x - 1)^2 + \ln \sqrt{3x - 1} - \ln x^7 - \ln(6 - 7x^2)^{\frac{3}{2}} \{ \text{as } \log ab = \log a + \log b \text{ \& } \log \frac{a}{b} = \log a - \log b \}$$

$$\Rightarrow \ln y = 2 \ln(x - 1) + \frac{1}{2} \ln(3x - 1) - 7 \ln x - \frac{3}{2} \ln(6 - 7x^2) \{ \text{as } \ln x^n = n \ln x \}$$

Differentiating both sides w.r.t, we have

$$\begin{aligned} \frac{1}{y} \frac{dy}{dx} &= \frac{2}{x-1} \frac{d(x-1)}{dx} + \frac{1}{2} \frac{1}{(3x-1)} \frac{d(3x-1)}{dx} - \frac{7}{x} - \frac{3}{2} \frac{1}{6-7x^2} \frac{d(6-7x^2)}{dx} \\ &= \frac{2}{x-1} + \frac{3}{2(3x-1)} - \frac{7}{x} + \frac{3(14x)}{2(6-7x^2)} \\ &= \frac{2}{x-1} + \frac{3}{2(3x-1)} - \frac{7}{x} + \frac{21x}{6-7x^2} \\ \Rightarrow \frac{dy}{dx} &= y \left[ \frac{2}{x-1} + \frac{3}{2(3x-1)} - \frac{7}{x} + \frac{21x}{6-7x^2} \right] \\ \Rightarrow \frac{dy}{dx} &= \frac{(x-1)^2 \sqrt{3x-1}}{x^7 (6-7x^2)^{\frac{3}{2}}} \left[ \frac{2}{x-1} + \frac{3}{2(3x-1)} - \frac{7}{x} + \frac{21x}{6-7x^2} \right] \end{aligned}$$

(Ans)

$$\Rightarrow \frac{1}{u} \frac{du}{dx} = \frac{d}{dx} (x \log (\ln x)) \text{ and } \frac{1}{v} \frac{dv}{dx} = \frac{d}{dx} (x \log (\sin^{-1} x))$$

$$\Rightarrow \frac{1}{u} \frac{du}{dx} = \left[ x \frac{1}{\ln x} \cdot \frac{1}{x} + 1 \cdot \log (\ln x) \right] \text{ and } \frac{1}{v} \frac{dv}{dx} = x \frac{1}{\sin^{-1} x} \cdot \frac{1}{\sqrt{1-x^2}} + 1 \cdot \log (\sin^{-1} x)$$

$$\Rightarrow \frac{du}{dx} = u \left[ \frac{1}{\ln x} + \log (\ln x) \right] \text{ and } \frac{dv}{dx} = v \left[ \frac{x}{\sin^{-1} x \sqrt{1-x^2}} + \log (\sin^{-1} x) \right]$$

$$\Rightarrow \frac{du}{dx} = (\ln x)^x \left[ \frac{1}{\ln x} + \log (\ln x) \right] \text{ and } \frac{dv}{dx} = (\sin^{-1} x)^x \left[ \frac{x}{\sin^{-1} x \sqrt{1-x^2}} + \log (\sin^{-1} x) \right]$$

Now,  $\frac{dy}{dx} = \frac{d}{dx} (u + v) = \frac{du}{dx} + \frac{dv}{dx}$

$$= (\ln x)^x \left[ \frac{1}{\ln x} + \log (\ln x) \right] + (\sin^{-1} x)^x \left[ \frac{x}{\sin^{-1} x \sqrt{1-x^2}} + \log (\sin^{-1} x) \right] \text{ (Ans)}$$

**Differentiation of parametric function**

Sometimes the variables x and y of a function is represent by function of another variable ‘t’, which is called as a parameter. Such type of representation of a fnction is called parametric form. For example equation of circle can be given by x = r cos t, y = r sin t.

Here x, y both are functions of parameter ‘t’.

So, this form of the function is called parametric form.

### Differentiation of a function w.r.t another function

---

#### Example-28

Find the derivative of  $\tan x$  w.r.t  $\cot x$  (2017-w)

Ans.

Let  $y = \tan x$  and  $z = \cot x$

$$\frac{dy}{dx} = \sec^2 x, \frac{dz}{dx} = -\operatorname{cosec}^2 x$$

$$\text{Now, } \frac{dy}{dz} = \frac{\left(\frac{dy}{dx}\right)}{\left(\frac{dz}{dx}\right)} = \frac{\sec^2 x}{-\operatorname{cosec}^2 x} = -\sec^2 x \sin^2 x \quad \text{, Hence } \frac{d(\tan x)}{d(\cot x)} = -\sec^2 x \sin^2 x$$

#### Example – 29

Find the derivative of  $e^{2 \log x}$  w.r.t  $2x^2$  [2018-S, 2017-w]

Ans.

$y = e^{2 \log x}$  and  $z = 2x^2$

$$\frac{dy}{dx} = \frac{d}{dx}(e^{2 \log x}) = \frac{d}{dx}(e^{\log x^2}) = \frac{d}{dx}(x^2) = 2x$$

$$\frac{dz}{dx} = \frac{d}{dx}(2x^2) = 4x$$

$$\text{Hence } \frac{dy}{dz} = \left(\frac{dy}{dx}\right) / \left(\frac{dz}{dx}\right) = \frac{2x}{4x} = \frac{1}{2}$$

### Differentiation of implicit function

---

Functions of the form  $F(x,y) = 0$  where  $x$  and  $y$  cannot be separated or in other words  $y$  cannot be expressed in terms of  $x$  is called Implicit function.

e.g.  $x^2 + y^2 - 25 = 0$

$$x^y = y^x$$

$$x^2 y + y^2 x + xy = 25 \quad \text{etc}$$

Derivative of Implicit functions can be found without expressing  $y$  explicitly in terms of  $x$ . Simply we differentiate both side w.r.t  $x$  and express  $\frac{dy}{dx}$  in terms of both  $x$  and  $y$ .

#### Example – 34

Find  $\frac{dy}{dx}$  when  $x^3 + y^3 - 3xy = 0$  [2015-S]

$dx$

**Ans.**

$$\text{Given } x^3 + y^3 - 3xy = 0$$

Differentiating both sides w.r.t x We have,

$$3x^2 + 3y^2 \frac{dy}{dx} - 3x \frac{dy}{dx} - 3 \cdot 1 \cdot y = 0$$

$$\frac{dy}{dx}(3y^2 - 3x) = 3y - 3x^2$$

$$\frac{dy}{dx} = \frac{3y - 3x^2}{3y^2 - 3x} = \frac{y - x^2}{y^2 - x} \quad (\text{Ans})$$

**Example – 36**

Find  $\frac{dy}{dx}$  if  $y^x = x^y$  [2014-S, 2016-S, 2017-w]

**Ans.** Given  $y^x = x^y$

Taking logarithm of both sides

$$\Rightarrow \ln y^x = \ln x^y$$

$$\Rightarrow x \ln y = y \ln x$$

Differentiating both sides w.r.t x, we have

$$\Rightarrow x \frac{1}{y} \frac{dy}{dx} + 1 \cdot \ln y = \frac{dy}{dx} \cdot \ln x + y \cdot \frac{1}{x}$$

$$\frac{x}{y} \frac{dy}{dx} + \ln y = \ln x \frac{dy}{dx} + \frac{y}{x}$$

$$\left(\frac{x}{y} - \ln x\right) \frac{dy}{dx} = \left(\frac{y}{x} - \ln y\right)$$

$$\Rightarrow \left(\frac{x - y \ln x}{y}\right) \frac{dy}{dx} = \frac{y - x \ln y}{x}$$

$$\Rightarrow \frac{dy}{dx} = \frac{y(y - x \ln y)}{x(x - y \ln x)}$$

$$\therefore \frac{dy}{dx} = \frac{y(y - x \ln y)}{x(x - y \ln x)}$$

**Example – 37**

Find  $\frac{dy}{dx}$  if  $y^2 \cot x = x^2 \cot y$

**Ans.**  $y^2 \cot x = x^2 \cot y$

Differentiating both sides w.r.t x,

$$\Rightarrow 2y \frac{dy}{dx} \cot x + y^2 (-\operatorname{cosec}^2 x) = 2x \cot y + x^2 (-\operatorname{cosec}^2 y \frac{dy}{dx})$$

$$\Rightarrow 2y \cot x \frac{dy}{dx} - y^2 \operatorname{cosec}^2 x = 2x \cot y - x^2 \operatorname{cosec}^2 y \frac{dy}{dx}$$

$$\Rightarrow (2y \cot x + x^2 \operatorname{cosec}^2 y) \frac{dy}{dx} = 2x \cot y + y^2 \operatorname{cosec}^2 x$$

$$\Rightarrow \frac{dy}{dx} = \frac{2x \cot y + y^2 \operatorname{cosec}^2 x}{2y \cot x + x^2 \operatorname{cosec}^2 y}$$

## Successive Differentiation

If  $f$  is a differential function of  $x$ , then the derivative of  $f(x)$  may be again differentiable w.r.t  $x$ . If  $f'(x) = \frac{df}{dx}$  then  $f'(x)$  is called first derivative of  $f$ .

If  $f'(x)$  is differentiable, and  $\frac{df'(x)}{dx} = f''(x)$ , then  $f''(x)$  is called the 2<sup>nd</sup> order derivative of  $f(x)$  w.r.t  $x$ .

The above process can be successively continued to obtain derivative functions of higher orders.

### Notations

1<sup>st</sup> Order derivatives  $\rightarrow \frac{dy}{dx}, y', y_1, Dy, f'(x)$

2<sup>nd</sup> Order derivatives  $\rightarrow \frac{d^2y}{dx^2}, y'', y_2, D^2y, f''(x)$

3<sup>rd</sup> Order derivatives  $\rightarrow \frac{d^3y}{dx^3}, y''', y_3, D^3y, f'''(x)$

.

.

.

$n^{\text{th}}$  Order derivatives  $\rightarrow \frac{d^ny}{dx^n}, y^{(n)}, y_n, D^ny, f^{(n)}(x)$

### Example – 1

Find 2<sup>nd</sup> order derivatives of following function.

(i)  $y = x^5 + 4x^3 - 2x^2 + 1$

(ii)  $y = \log_e x$

(iii)  $y = \sqrt{x^2 + 1}$

(iv)  $y = \frac{1}{\sqrt{x}}$

(i)  $y' = \frac{dy}{dx} = \frac{d}{dx}(x^5 + 4x^3 - 2x^2 + 1) = 5x^4 + 12x^2 - 4x + 0$   
 $= 5x^4 + 12x^2 - 4x$

$$y_2 = \frac{d^2y}{dx^2} = \frac{d}{dx} \left( \frac{dy}{dx} \right) = \frac{d}{dx} (5x^4 + 12x^2 - 4x) = 20x^3 + 24x - 4 \quad (\text{Ans})$$

$$(ii) \quad y_1 = \frac{d}{dx} (\log_e x) = \frac{1}{x}$$

$$(iii) \quad y_2 = \frac{dy_1}{dx} = \frac{d\left(\frac{1}{x}\right)}{dx} = -\frac{1}{x^2} \quad (\text{ans})$$

(iii)

$$y_1 = \frac{1}{2\sqrt{x^2+1}} \frac{d}{dx} (x^2 + 1) \quad (\text{Chain Rule})$$

$$= \frac{2x+0}{2\sqrt{x^2+1}} = \frac{x}{\sqrt{x^2+1}}$$

$$y_2 = \frac{dy_1}{dx} = \frac{d\left(\frac{x}{\sqrt{x^2+1}}\right)}{dx}$$

$$= \frac{1 \cdot \sqrt{x^2+1} - x \cdot \frac{1}{2\sqrt{x^2+1}} \frac{d(x^2+1)}{dx}}{(\sqrt{x^2+1})^2} \quad (\text{applying division formula of derivative})$$

$$= \frac{\sqrt{(x^2+1)} - \frac{x}{2\sqrt{x^2+1}} \cdot 2x}{x^2+1}$$

$$= \frac{(x^2+1) - x^2}{\sqrt{x^2+1} (x^2+1)} = \frac{1}{(x^2+1)^{3/2}} \quad (\text{Ans})$$

$$(iv) \quad y_1 = \frac{d}{dx} \left( \frac{1}{\sqrt{x}} \right) = \frac{d}{dx} (x^{-1/2}) = -\frac{1}{2} x^{-1/2-1} = -\frac{1}{2} x^{-3/2}$$

$$y_2 = \frac{dy_1}{dx} = -\frac{1}{2} \left( -\frac{3}{2} \right) x^{-3/2-1} = \frac{3}{4} x^{-5/2} = \frac{3}{4x^{5/2}}$$

### Example – 2

Find  $y_1$  and  $y_2$  if  $y = \log(\sin x)$  (2018-S) **Ans.**

$$y_1 = \frac{d}{dx} \log(\sin x) = \frac{1}{\sin x} \cos x = \cot x$$

$$y_2 = \frac{dy_1}{dx} = \frac{d}{dx} (\cot x) = -\operatorname{cosec}^2 x \quad (\text{Ans})$$

### Example – 3

If  $x = at^2$ ,  $y = 2at$  then find  $\frac{d^2y}{dx^2}$  **Ans.**

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)} = \frac{\frac{d}{dt}(2at)}{\frac{d}{dt}(at^2)} = \frac{2a}{2at} = \frac{1}{t}$$

$$\text{Now } \frac{d^2y}{dx^2} = \frac{d}{dx} \left( \frac{dy}{dx} \right) = \frac{d}{dt} \left( \frac{dy}{dx} \right) \cdot \frac{dt}{dx}$$

$$= \frac{\frac{d}{dt}\left(\frac{1}{t}\right)}{\frac{dx}{dt}} = \frac{-\frac{1}{t^2}}{2at} = -\frac{1}{2at^3}$$

## Partial Differentiation

The functions studied so far are of a single independent variable. There are functions which depends on two or more variables. Example, the pressure(P) of a given mass of gas is dependent on its volume(v) and temperature (T).

### Functions of two variable

A function  $f : X \times Y$  to  $Z$  is a function of two variables if there exist a unique element  $z = f(x,y)$  in  $Z$  corresponding to every pair  $(x,y)$  in  $X \times Y$ .

Domain of  $f$  is  $X \times Y$ .

$f(X \times Y)$  is the range of  $f$ .  $\{ f(X \times Y) \subset Z \}$

**Notation :** -  $z = f(x,y)$  means  $z$  is a function of two variables  $x$  and  $y$ .

### Limit of a function of two variable

A function  $f(x,y)$  tends to limit  $l$  as  $(x,y) \rightarrow (a,b)$ , .If given  $\epsilon > 0$ , there exist  $\delta > 0$  such that  $|f(x,y) - l| < \epsilon$  whenever  $0 < |(x,y) - (a,b)| < \delta$ .

### Continuity

A function  $f(x,y)$  is said to be continuous at a point  $(a,b)$  if

- (i)  $f(a,b)$  is defined
- (ii)  $\lim_{(x,y) \rightarrow (a,b)} f(x,y)$  exists.
- (iii)  $\lim_{(x,y) \rightarrow (a,b)} f(x,y) = f(a,b)$

Finding limits and testing continuity of functions of two variable is beyond our syllabus so we have to skip these topics here.

### Partial derivatives

Let  $z = f(x,y)$  be function of two variables.

If variable  $x$  undergoes a change  $\delta x$ , while  $y$  remains constant, then  $z$  undergoes a change written as  $\delta z$

Now,  $\delta z = f(x+\delta x, y) - f(x, y)$

If  $\frac{\delta z}{\delta x}$  exist as  $\delta x \rightarrow 0$ , then we write the partial derivative of  $z$  w.r.t  $x$  as



$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial x} = f_x = z_x = \text{_____}$$

Similarly partial derivative of z w.r.t y ,

$$\frac{\partial z}{\partial y} = \frac{\partial z}{\partial y} = f_y = z_y = \text{_____}$$

$\frac{\partial}{\partial x}$  ,  $\frac{\partial}{\partial y}$  symbols are used to notify the partial differentiation.

### Note

As from above theory it is clear when partial differentiation w.r.t x is taken , then y is treated as constant and vice – versa. (All the formulae and techniques used in derivative chapter remain same here)

## 2<sup>nd</sup> Order Partial Differentiation

If we differentiate the  $\frac{\partial z}{\partial x}$  ,  $\frac{\partial z}{\partial y}$  w.r.t x or y , then we set higher order partial derivatives as follows.

1<sup>st</sup> Order Partial Derivatives  $\frac{\partial z}{\partial x}$  ,  $\frac{\partial z}{\partial y}$ .

2<sup>nd</sup> Order Partial Derivatives

$$\frac{\partial^2 z}{\partial x^2} = \frac{\partial}{\partial x} \left( \frac{\partial z}{\partial x} \right) = z_{xx} = f_{xx}$$

$$\frac{\partial^2 z}{\partial y \partial x} = \frac{\partial}{\partial y} \left( \frac{\partial z}{\partial x} \right) = z_{yx} = f_{yx}$$

$$\frac{\partial^2 z}{\partial x \partial y} = \frac{\partial}{\partial x} \left( \frac{\partial z}{\partial y} \right) = z_{xy} = f_{xy}$$

$$\frac{\partial^2 z}{\partial y^2} = \frac{\partial}{\partial y} \left( \frac{\partial z}{\partial y} \right) = z_{yy} = f_{yy}$$

Note:  $f_{yx} = f_{xy}$  when partial derivatives are continuous .

### Example -1

Find  $\frac{\partial z}{\partial x}$  ,  $\frac{\partial z}{\partial y}$

(i)  $z = 2x^2y + xy^2 + 5xy.$

$$(ii) \quad z = \tan^{-1}\left(\frac{x}{y}\right)$$

**Ans.**

$$(i) \quad z = 2x^2y + xy^2 + 5xy$$

$$\frac{\partial z}{\partial x} = \frac{\partial}{\partial x}(2x^2y) + \frac{\partial}{\partial x}(xy^2) + \frac{\partial}{\partial x}(5xy) \quad (\text{Here } y \text{ is treated as constant})$$

$$= 2y \frac{\partial}{\partial x}(x^2) + y^2 \frac{\partial}{\partial x}(x) + 5y \frac{\partial}{\partial x}(x)$$

$$= 2y \cdot 2x + y^2 \cdot 1 + 5y \cdot 1$$

$$= 4xy + y^2 + 5y.$$

$$\frac{\partial z}{\partial y} = 2x^2 \frac{\partial y}{\partial y} + x \frac{\partial y^2}{\partial y} + 5x \frac{\partial y}{\partial y}$$

$$= 2x^2 + x \cdot 2y + 5x = 2x^2 + 2xy + 5x$$

$$(ii) \quad z = \tan^{-1}\left(\frac{x}{y}\right)$$

$$\frac{\delta z}{\delta x} = \frac{1}{1+\left(\frac{x}{y}\right)^2} \frac{\partial}{\partial x} \left(\frac{x}{y}\right) = \frac{1}{\frac{y^2+x^2}{y^2}} \cdot \frac{1}{y}$$

$$= \frac{y^2}{y(x^2+y^2)} = \frac{y}{x^2+y^2}$$

$$\frac{\partial z}{\partial y} = \frac{1}{1+\left(\frac{x}{y}\right)^2} \frac{\partial}{\partial y} \left(\frac{x}{y}\right) = \frac{1}{\frac{y^2+x^2}{y^2}} \left(\frac{-x}{y^2}\right)$$

$$= \frac{-x}{x^2+y^2}$$

### **Homogenous function and Euler's theorem**

---

#### **Homogenous function**

A function  $f(x, y)$  is said to be homogenous in  $x$  and  $y$  of degree  $n$  iff  $(tx, ty) = t^n f(x, y)$  where  $t$  is any constant.

#### **Example – 4**

Test whether the following functions are homogenous or not. If homogenous then find their degree.

$$(i) \quad 2xy^2 + 3x^2y \qquad (ii) \quad \sin^{-1}\left(\frac{x}{y}\right)$$

**Ans.**

$$(i) \quad \text{Let } f(x, y) = 2xy^2 + 3x^2y \quad f(tx, ty) = 2(tx)(ty)^2 + 3(tx)^2(ty) = 2txt^2y^2 + 3t^2x^2ty \\ = t^3(2xy^2 + 3x^2y) = t^3f(x, y)$$

Hence  $f(x, y)$  is a homogenous function of degree 3.

- (ii) Let  $f(x, y) = \sin^{-1}\left(\frac{x}{y}\right)$   
 $f(tx, ty) = \sin^{-1}\left(\frac{tx}{ty}\right) = \sin^{-1}\frac{x}{y} = t^0 \sin^{-1}\left(\frac{x}{y}\right) = t^0 f(x, y)$   
 Hence  $f(x, y)$  is a homogenous function of degree '0'.

**Note**

- (i) If each term in the expression of a function is of the same degree then the function is homogenous.  
 (ii) If  $z$  is a homogenous function of  $x$  and  $y$  of degree  $n$ , then  $\frac{\partial z}{\partial x}$  and  $\frac{\partial z}{\partial y}$  are also homogenous of degree  $n-1$ .  
 (iii) If  $z = f(x, y)$  is a homogenous function of degree  $n$ , then we can write it as  $z = x^n \Phi\left(\frac{y}{x}\right)$

e.g. In example -4(i)  $2x y^2 + 3x^2 y$  is homogenous function of degree 3.  
 Now  $f(x, y) = 2xy^2 + 3x^2 y = x^3 \left(2\left(\frac{y}{x}\right)^2 + 3\left(\frac{y}{x}\right)\right) = x^3 \Phi\left(\frac{y}{x}\right)$

Similarly in Example - 4 (iii),  $f(x,y)$  is of degree 1.

$$\text{Now } f(x, y) = \frac{3x^2+2y^2}{x+y} = \frac{x^2}{x} \left(\frac{3+2\left(\frac{y}{x}\right)^2}{1+\frac{y}{x}}\right) = x \left(\frac{3+2\left(\frac{y}{x}\right)^2}{1+\frac{y}{x}}\right) = x \Phi\left(\frac{y}{x}\right)$$

**Euler's theorem**

If  $z$  is a homogenous function of degree  $n$ , then  $x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = n z$ . [2014-S]

**Proof: -**

Since  $z$  is a homogenous function of degree  $n$ , so  $z$  can be written as

$$z = x^n \Phi\left(\frac{y}{x}\right)$$

$$\begin{aligned} \text{Now } \frac{\partial z}{\partial x} &= n x^{n-1} \Phi\left(\frac{y}{x}\right) + x^n \Phi'\left(\frac{y}{x}\right) \cdot \frac{\partial}{\partial x}\left(\frac{y}{x}\right) \\ &= n x^{n-1} \Phi\left(\frac{y}{x}\right) + x^n \Phi'\left(\frac{y}{x}\right) \left(-\frac{y}{x^2}\right) \\ &= n x^{n-1} \Phi\left(\frac{y}{x}\right) - x^{n-2} y \Phi'\left(\frac{y}{x}\right) \text{----- (1)} \end{aligned}$$

$$\begin{aligned} \text{Similarly, } \frac{\partial z}{\partial y} &= x^n \Phi'\left(\frac{y}{x}\right) \frac{\partial}{\partial y}\left(\frac{y}{x}\right) \\ &= x^n \Phi'\left(\frac{y}{x}\right) \left(\frac{1}{x}\right) = x^{n-1} \Phi'\left(\frac{y}{x}\right) \text{----- (2)} \end{aligned}$$

Now  $x \times$  Equation (1) +  $y \times$  Equation (2)

$$\Rightarrow x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = x \{n x^{n-1} \Phi\left(\frac{y}{x}\right) - x^{n-2} y \Phi'\left(\frac{y}{x}\right)\} + y x^{n-1} \Phi'\left(\frac{y}{x}\right)$$

$$= n x^n \phi\left(\frac{y}{x}\right) - x^{n-1} y \phi'\left(\frac{y}{x}\right) + x^{n-1} y \phi'\left(\frac{y}{x}\right)$$

$$= n x^n \phi\left(\frac{y}{x}\right) = n z \text{ (proved)}$$

**Example – 5**

Verify Euler's theorem for  $z = \frac{y}{x}$  [2014-S]

**Ans.**  $\frac{\partial z}{\partial x} = \frac{\partial}{\partial x} \left(\frac{y}{x}\right) = -\frac{y}{x^2}$

$$\frac{\partial z}{\partial y} = \frac{\partial}{\partial y} \left(\frac{y}{x}\right) = \frac{1}{x}$$

Here  $z = f(x, y) = \frac{y}{x}$

$$F(tx, ty) = \frac{ty}{tx} = \frac{y}{x} = f(x, y)$$

Hence  $f(x, y)$  is a homogenous function of degree 0.

Statement of Euler's theorem is  $x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = n z$  (here  $n=0$ )

$$\Rightarrow x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = 0 \cdot z = 0$$

Now we have to verify it.

From above

$$\text{L.H.S} = x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = x \left(-\frac{y}{x^2}\right) + y \cdot \frac{1}{x} = -\frac{y}{x} + \frac{y}{x} = 0 = \text{R.H.S}$$

Hence Euler's theorem is verified.

**Exercise****Question with short answers (2 marks)**

1) if  $z = \sin \frac{x}{y}$  find  $\frac{\partial z}{\partial x}$  and  $\frac{\partial z}{\partial y}$ .

2) If  $f(x,y) = \sqrt{x^2 + y^2}$ , find  $f_x, f_y$ .

3) If  $f(x,y) = \log(x^2 + y^2 - 2xy)$  find  $f_{xx}, f_{yx}, f_{xy}$

**Questions with long answers (5 marks)**

4) Given  $f(u,v) = \frac{2u-3v}{u^2+v^2}$ , find  $f_u(2,1)$  and  $f_v(2,1)$

# UNIT-IV

## INTEGRATION

### Introduction

---

Calculus deals with some important geometrical problem related to draw a tangent of a curve and determine area of a region under a curve. In order to solve these problems we use differentiation and integration respectively.

In the previous lesson, we have studied derivative of a function. After studying differentiation it is natural to study the inverse process called integration.

### Objectives

---

- After completion of this topic you will able to
1. Explain integration as inverse process of differentiation.
  2. State types of integration.
  3. State integral of some standard functions like  $x^n$ ,  $\sin x$ ,  $\cos x$ , ...,  $\sin^{-1} x$ , .....  $a^x$  etc.
  4. State properties of integration.
  5. Find integration of algebraic, trigonometric, inverse trigonometric functions using standard integration formulae.
  6. Evaluate different integrals by applying substitution method and integration by parts method.

### Expected Background Knowledge

---

1. Trigonometry
2. Derivative

### Integration (Primitive or Anti derivative)

---

Integration is the reverse process of differentiation.

If  $\frac{df(x)}{dx} = g(x)$ , then the integration of  $g(x)$  w.r.t  $x$  is  $\int g(x)dx = f(x) + c$

- ➔ The Symbol  $\int$  is used to denote the operation of integration called as Integral sign.  $\square$  The function (here  $g(x)$ ) is called the integrand.
- ➔ 'dx' denote that the Integration is to be performed w.r.t  $x$  ( $x$  is the variable of Integration).
- ➔ 'c' is the constant of Integration (which gives family of curves)

➔ Integrate means to find the Integral of the function and the process is known as Integration

### Types of Integration

---

Integration are of two types:- i) Indefinite ii) definite The integration

written in the form  $\int g(x)dx$  is called indefinite integral.

The integration written in the form  $\int_a^b g(x)dx$  is called definite integral.

In this chapter we only discuss the indefinite integrals. The definite integrals will be discussed in the next chapter.

### Algebra of Integrals

---

$$i. \quad \int [f(x) \pm g(x)]dx = \int f(x)dx \pm \int g(x)dx$$

$$ii. \quad \int \lambda f(x)dx = \lambda \int f(x)dx \quad \text{for any constant}$$

$\lambda.$

$$iii. \quad \frac{d}{dx} (\lambda \int f(x)dx) = \lambda \frac{d}{dx} (\int f(x)dx) = \lambda f(x)$$

### Simple Integration Formula of some standard functions

---

$$i) \quad \int k dx = kx + c$$

$$ii) \quad \int x^n dx = \frac{x^{n+1}}{n+1} + c, \quad n \neq -1$$

$$iii) \quad \int \frac{1}{x} dx = \ln|x| + c$$

$$iv) \quad \int a^x dx = \frac{a^x}{\ln a} + c$$

$$v) \quad \int e^x dx = e^x + c$$

$$vi) \quad \int \sin x dx = -\cos x + c \quad vii)$$

$$\int \cos x dx = \sin x + c \quad vii)i)$$

$$\int \sec^2 x dx = \tan x + c \quad ix)$$

$$\int \operatorname{cosec}^2 x dx = -\cot x + c$$

$$x) \quad \int \sec x \tan x dx = \sec x + c$$

$$xi) \quad \int \operatorname{cosec} x \cot x dx = -\operatorname{cosec} x + c$$

$$xii) \quad \int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1} x + c$$

$$\text{xii) } \int \frac{-1}{\sqrt{1-x^2}} dx = \cos^{-1}x + c$$

$$\text{xiv) } \int \frac{1}{1+x^2} dx = \tan^{-1}x + c \quad \text{xv) } \int \frac{-1}{1+x^2} dx = \cot^{-1}x + c \quad \text{xvi) }$$

$$\int \frac{1}{x\sqrt{x^2-1}} dx = \sec^{-1}x + c \quad \text{xvii) } \int \frac{-1}{x\sqrt{x^2-1}} dx = \operatorname{cosec}^{-1}x + c$$

### Methods of integration

1. Integration by using standard formula.
2. Integration by substitution.
3. Integration by parts.

### 1. INTEGRATION BY USING FORMULAS:-

**Example -1** Evaluate the following

$$\text{(i) } \int (5x^3 + 2x^5 - 7x + \frac{1}{\sqrt{x}} + \frac{5}{x}) dx$$

$$\text{Ans : } \int (5x^3 + 2x^5 - 7x + \frac{1}{\sqrt{x}} + \frac{5}{x}) dx$$

$$= 5 \int x^3 dx + 2 \int x^5 dx - 7 \int x dx + \int x^{-1/2} dx + 5 \int \frac{dx}{x} \{ \text{by algebra of integration} \}$$

$$= 5 \times \frac{x^{3+1}}{3+1} + 2 \times \frac{x^{5+1}}{5+1} - 7 \times \frac{x^{1+1}}{1+1} + \frac{x^{-1/2+1}}{-1/2+1} + 5 \ln|x| + c$$

$$= 5 \times \frac{x^4}{4} + 2 \times \frac{x^6}{6} - 7 \times \frac{x^2}{2} + \frac{x^{1/2}}{1/2} + 5 \ln|x| + c$$

2

$$= \frac{5x^4}{4} + \frac{x^6}{3} - \frac{7x^2}{2} + 2x^{1/2} + 5 \ln x + c$$

$$= \frac{5x^4}{4} + \frac{x^6}{3} - \frac{7x^2}{2} + 2\sqrt{x} + 5 \ln x + c$$

$$\text{(ii) } \int \left( \frac{3x^4 - 5x^3 + 4x^2 - x + 2}{x^3} \right) dx \quad \text{Ans :}$$

$$= \int \left( \frac{3x^4 - 5x^3 + 4x^2 - x + 2}{x^3} \right) dx$$

$$= \int \frac{3x^4}{x^3} dx - \int \frac{5x^3}{x^3} dx + \int \frac{4x^2}{x^3} dx - \int \frac{x}{x^3} dx + \int \frac{2}{x^3} dx$$

$$= 3 \int x dx - 5 \int dx + 4 \int \frac{dx}{x} - \int x^{-2} dx + 2 \int x^{-3} dx$$

$$= 3 \times \frac{x^{1+1}}{1+1} - 5x + 4 \ln x - \frac{x^{-2+1}}{-2+1} + \frac{2x^{-3+1}}{-3+1} + c$$

$$= 3 \times \frac{x^2}{2} - 5x + 4 \ln x + \frac{1}{x} - \frac{1}{x^2} + c$$

$$= \frac{3x^2}{2} - 5x + 4\ln x + \frac{1}{x} - \frac{1}{x^2} + c$$

## 2. INTEGRATION BY SUBSTITUTION:-

When the integral  $\int f(x)dx$  cannot be determined by the standard formulae then we may reduce it to another form by changing the independent variable 'x' by another variable t (as  $x=\phi(t)$ ) which can be integrated easily. This is called substitution method.

$$\int f(x)dx = \int f(x)\frac{dx}{dt}dt = \int f[\phi(t)]\phi'(t)dt, \quad \text{where } x=\phi(t).$$

The substitution  $x=\phi(t)$  depends upon the nature of the given integral and has to be properly chosen so that integration is easier after substitution. The following types of substitution are very often used in Integrations.

### TYPE – I

$$\int f(ax + b)dx \Rightarrow dx = \frac{1}{a} dt$$

$$\text{Put } ax + b = t$$

$$\begin{aligned} a dx &= dt \\ \therefore \int f(ax + b)dx &= \int f(t)\frac{1}{a}dt = \frac{1}{a} \int f(t)dt \end{aligned}$$

### TYPE – II

$$\int x^{n-1} f(x^n)dx$$

$$\text{Put } x^n = t$$

$$nx^{n-1}dx = dt$$

$$\Rightarrow x^{n-1}dx = \frac{dt}{n}$$

$$\therefore \int x^{n-1} f(x^n)dx = \int f(t)\frac{dt}{n} = \frac{1}{n} \int f(t)dt$$

### TYPE – III

$$\int \{f(x)\}^n \cdot f'(x)dx$$

$$\text{Put } f(x)=t$$

Differentiate both sides w.r.t x,

$$f'(x) = \frac{dt}{dx}$$

$$\Rightarrow \int \{f(x)\}^n \cdot f'(x)dx = \int t^n dt = \frac{t^{n+1}}{n+1} + c$$



$$= \frac{[f(x)]^{n+1}}{n+1} + c \quad (\because t = f(x))$$

**TYPE-IV**

$$\int \frac{f'(x)}{f(x)} dx$$

Put  $f(x)=t$

$$\Rightarrow f'(x)dx = dt$$

$$\therefore \int \frac{f'(x)}{f(x)} dx = \int \frac{dt}{t} = \ln|t| + c = \ln|f(x)| + c \quad (\because f(x) = t)$$

**SOME USE FULL RESULTS**

$$1. \int \frac{dx}{ax+b}$$

**Ans :-** Put  $ax+b = t$

Differentiate both sides w.r.t  $x$ ,

$$a = \frac{dt}{dx}$$

$$\Rightarrow dx = \frac{dt}{a}$$

$$\therefore \int \frac{dx}{ax+b} = \int \frac{dt/a}{t} = \frac{1}{a} \int \frac{dt}{t} = \frac{1}{a} \ln|t| = \frac{1}{a} \ln|ax+b| + c.$$

$\int \frac{dx}{ax+b} = \frac{1}{a} \ln ax+b  + c$
--

$$2. \int \cot x dx$$

**Ans:-**  $\int \cot x dx$

$$= \int \frac{\cos x}{\sin x} dx$$

Put  $\sin x = t$

Differentiate both sides w.r.t  $x$ ,

$$\cos x = \frac{dt}{dx}$$

$$\Rightarrow dt = \cos x dx$$

$$\therefore \int \frac{\cos x dx}{\sin x} = \int \frac{dt}{t} = \ln|t| = \ln|\sin x| + c$$

### INTEGRATION OF SOME TRIGONOMETRIC FUNCTIONS

#### Example – 3

i) Evaluate  $\int \sin 3x \cos 2x dx$

Ans :-  $\int \sin 3x \cos 2x dx$

$$\begin{aligned} &= \frac{1}{2} \int \sin(3x + 2x) + \sin(3x - 2x) dx \\ &= \frac{1}{2} \int (\sin 5x + \sin x) dx \\ &= \frac{1}{2} \int \sin 5x dx + \frac{1}{2} \int \sin x dx \\ &= \frac{1}{2} \times \frac{-\cos 5x}{5} + \frac{1}{2} (-\cos x) + c \\ &= \frac{-1}{10} \cos 5x - \frac{1}{2} \cos x + c \\ &= \frac{-1}{10} (\cos 5x - 5 \cos x) + c \end{aligned}$$

ii) Evaluate  $\int \sin 2x \sin x dx$

Ans :-  $\int \sin 2x \sin x dx$

$$\begin{aligned} &= \frac{1}{2} \int \cos(2x - x) - \cos(2x + x) dx \\ &= \frac{1}{2} \int (\cos x - \cos 3x) dx \\ &= \frac{1}{2} \int \cos x dx - \frac{1}{2} \int \cos 3x dx \\ &= \frac{1}{2} \times \sin x - \frac{1}{2} \times \frac{\sin 3x}{3} + c \\ &= \frac{1}{2} \sin x - \frac{1}{6} \sin 3x + c = \frac{1}{6} (3 \sin x - \sin 3x) + c \end{aligned}$$

### INTEGRATION BY TRIGONOMETRIC SUBSTITUTION

#### TRIGONOMETRIC IDENTITIES

$$1 - \sin^2 \theta = \cos^2 \theta \text{ (or } 1 - \cos^2 \theta = \sin^2 \theta)$$

$$\tan^2 \theta + 1 = \sec^2 \theta \text{ (also } \cot^2 \theta + 1 = \operatorname{cosec}^2 \theta)$$

$$\sec^2 \theta - 1 = \tan^2 \theta \text{ (also } \operatorname{cosec}^2 \theta - 1 = \cot^2 \theta)$$

□ the integrand of the form  $\sqrt{a^2 - x^2}, \sqrt{x^2 + a^2}, \sqrt{x^2 - a^2}$  can be simplified by putting

$X=a \sin \theta$
$X=a \tan \theta$
$X=a \sec \theta$
$X=a \cos \theta$
$X=a \cot \theta$
$X=a \operatorname{cosec} \theta$

**Note**

1. The integrand of the form  $a^2 - x^2$  can be simplify by putting  $x = a \sin \theta$  (or  $x = a \cos \theta$ )
2. The integrand of the form  $x^2 + a^2$  can be simplify by putting  $x = a \tan \theta$  (or  $x = a \cot \theta$ )
3. The integrand of the form  $x^2 - a^2$  can be simplify by putting  $x = a \sec \theta$  (or  $x = a \operatorname{cosec} \theta$ )

**Example -4**

i) Integrate  $\int \frac{dx}{\sqrt{a^2 - x^2}}$

Ans :-  $\int \frac{dx}{\sqrt{a^2 - x^2}}$

Let  $x = a \sin \theta$

Differentiate both sides w.r.t x

$$dx = a \cos \theta d\theta$$

And  $x = a \sin \theta \Rightarrow \theta = \sin^{-1} \frac{x}{a}$

Hence  $\int \frac{dx}{\sqrt{a^2 - x^2}} = \int \frac{a \cos \theta d\theta}{\sqrt{a^2 - a^2 \sin^2 \theta}} = \int \frac{a \cos \theta}{a \cos \theta} d\theta = \int d\theta = \theta + c = \sin^{-1} \frac{x}{a} + c$

$\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{x}{a} + c$
--

ii) Integrate  $\int \frac{dx}{x^2 + a^2}$

Ans :-  $\int \frac{dx}{x^2 + a^2}$

Let  $x = a \tan \theta$

differentiating both sides w.r.t x,

$$dx = a \sec^2 \theta d\theta$$

And  $x = \tan \theta$

$$\Rightarrow \boxed{\theta = \tan^{-1} \frac{x}{a}}$$

$$\begin{aligned} \text{Hence } \int \frac{dx}{x^2+a^2} &= \int \frac{a \sec^2 \theta d\theta}{a^2 \tan^2 \theta + a^2} \\ &= \int \frac{a \sec^2 \theta d\theta}{a^2 (\tan^2 \theta + 1)} \\ &= \int \frac{\sec^2 \theta d\theta}{a \sec^2 \theta} \\ &= \frac{1}{a} \int d\theta = \frac{1}{a} \theta + c = \frac{1}{a} \tan^{-1} \frac{x}{a} + c \end{aligned}$$

$$\boxed{\int \frac{dx}{x^2+a^2} = \frac{1}{a} \tan^{-1} \frac{x}{a} + c}$$

### 3. INTEGRATION BY PARTS:-

If v & w are two differentiation function of x, then

$$\frac{d}{dx}(vw) = v \frac{dw}{dx} + w \frac{dv}{dx}$$

$$\text{Or } v \frac{dw}{dx} = \frac{d}{dx}(vw) - w \frac{dv}{dx}$$

Integrating both sides,

$$\begin{aligned} \int v \frac{dw}{dx} dx &= \int \frac{d}{dx}(vw) dx - \int w \frac{dv}{dx} dx \\ &= vw - \int w \frac{dv}{dx} dx \end{aligned}$$

$$\text{Let } u = \frac{dv}{dx} \text{ then } w = \int u dx$$

Then the above result can be written as  $\int uv dx = (\int u dx) v - \int (\int u dx) \times \frac{dv}{dx} dx$ .

This rule is called integration by parts and is used to integrate the product of two functions

#### Integration of the product of two functions

$$= (\text{integral of first function}) \times \text{second function} - \text{integral of } (\text{integral of first} \times \text{derivative of second})$$

$$\boxed{\text{Int. of product} = (\text{int. first}) \times \text{second} - \int (\text{int first} \times \text{der second}) dx}$$

- Before applying integration by parts we have follow some important things which are listed below.
1. In above formula there are two functions one is u and other one is v. The function 'u' is called the 1<sup>st</sup> function where as 'v' is called as the 2<sup>nd</sup> function.
  2. The choice of 1<sup>st</sup> function is made basing on the order ETALI . The meaning of these letters is given below.

E – Exponential function

T – Trigonometric function

A – Algebraic function

L – Logarithmic function

I – inverse trigonometric function

The following table-1 gives a proper choice of 1<sup>st</sup> and 2<sup>nd</sup> function in certain cases. Here  $m \in \mathbb{N}$ , n may be zero or any positive integer.

**Table-1**

Function to be integrated	first function	second function
$x^n e^x$	$e^x$	$x^n$
$x^n \sin x$	$\sin x$	$x^n$
$x^n \cos x$	$\cos x$	$x^n$
$x^n (\ln x)^m$	$x^n$	$(\ln x)^m$
$x^n \sin^{-1} x$	$x^n$	$\sin^{-1} x$
$x^n \cos^{-1} x$	$x^n$	$\cos^{-1} x$
$x^n \tan^{-1} x$	$x^n$	$\tan^{-1} x$

**Example – 11**

Integrate  $\int x \cos x \, dx$

**Ans :-**  $\int x \cos x \, dx$  { from table-1, 1<sup>st</sup> function =  $\cos x$  and 2<sup>nd</sup> function =  $x$  }

$$\begin{aligned}
 &= \{ \int (\cos x \, dx) \} x - \int (\int \cos x \, dx) \times \frac{dx}{dx} \cdot dx \\
 &= x \sin x - \int \sin x \cdot 1 \cdot dx \\
 &= x \sin x + \cos x + c
 \end{aligned}$$

**Example – 12**

Integrate  $\int x^2 e^x dx$

**Ans:-**  $\int x^2 e^x dx$  { 1<sup>st</sup> function =  $e^x$  and 2<sup>nd</sup> function =  $x^2$  }

$$= (\int e^x dx) \cdot x^2 - \int (\int e^x dx) \frac{d}{dx} (x^2) dx$$

$$= x^2 e^x - \int e^x \times 2x dx$$

$$= x^2 e^x - 2 \int x e^x dx \text{ { again by parts is applied taking } } e^x \text{ as 1<sup>st</sup> and } x \text{ as 2<sup>nd</sup> function. }$$

$$= x^2 e^x - 2 [ (\int e^x dx) \cdot x - \int (\int e^x dx) \cdot 1 \cdot dx ]$$

$$= x^2 e^x - 2x e^x + 2e^x + c = (x^2 - 2x + 2) e^x + c$$

**Exercise****1. Evaluate the following Integrals (2 marks questions)**

i)  $\int \frac{1}{x\sqrt{x}} dx$

ii)  $\int (x^{\frac{4}{7}} + \frac{1}{x^{\frac{1}{3}}}) dx$

iii)  $\int \frac{1 - \sin^3 x}{\sin^2 x} dx$

**2. Evaluate the following (2 marks questions)**

i)  $\int \frac{x^2 dx}{(1 + x^3)^2}$

ii)  $\int \sec^2 (3x + 5) dx$

iii)  $\int \frac{(\tan^{-1} x)^3}{1 + x^2} dx$

## Definite integral

### Introduction

---

It was stated earlier that integral can be considered as process of summation. In such case the integral is called definite integral.

### Objective

---

After completion of the topic you will be able to

1. Define and interpret geometrically the definite integral as a limit of sum.
2. State fundamental theorem of integral calculus.
3. State properties of definite integral.
4. Find the definite integral of some functions using properties.
5. Apply definite integral to find the area under a curve

### Expected Background knowledge

---

1. Functional value of a function at a point.
2. Integration.

### Definite Integral

---

Integration can be considered as a process of summation. In this case the integral is called as definite integrals.




Fig-1

**Definition:-**

Let  $f(x)$  be a continuous function in  $[a,b]$  as shown in Fig-1 . Divide  $[a,b]$  into  $n$  sub-intervals of length  $h_1, h_2, \dots, h_n$ . i.e.  $h_1 = x_1 - x_0, h_2 = x_2 - x_1, \dots, h_n = x_n - x_{n-1}$

Let  $v_r$  be any point in  $[x_{r-1}, x_r]$  i.e.  $v_1 \in [x_0, x_1], v_2 \in [x_1, x_2], \dots, v_n \in [x_{n-1}, x_n]$ .

Then the sum of area of the rectangles (as shown in fig) when  $n \rightarrow \infty$  is defined as the definite integral of  $f(x)$  from  $a$  to  $b$ , denoted by  $\int_a^b f(x) dx$

Here,  $a = \text{lower limit of integration}$   
 $b = \text{upper limit of integration}$

Mathematically,

$$\int_a^b f(x) dx = [h_1 f(v_1) + h_2 f(v_2) + \dots + h_n f(v_n)]$$

**Fundamental Theorem of Integral Calculus**

---

If  $f(x)$  is a continuous function in  $[a,b]$  and  $\int f(x) dx = \Phi(x) + c$ , then

$$\int_a^b f(x) dx = \Phi(b) - \Phi(a)$$

Note :- No arbitrary constants are used in definite integral.

**Example:**

1. Find  $\int_1^2 x^3 dx$

Ans.

First find  $\int x^3 dx = \frac{x^4}{4} + c$

Here,  $f(x) = x^3, \Phi(x) = \frac{x^4}{4}$

By fundamental theorem

$$\begin{aligned} \int_1^2 x^3 dx &= \Phi(2) - \Phi(1) \\ &= \frac{2^4}{4} - \frac{1^4}{4} = \frac{16}{4} - \frac{1}{4} = \frac{15}{4} \end{aligned}$$



2. Find  $\int_0^1 \frac{dx}{1+x^2}$

Ans.

$$\begin{aligned} \int_0^1 \frac{dx}{1+x^2} &= [\tan^{-1} x]_0^1 \\ &= \tan^{-1} 1 - \tan^{-1} 0 \\ &= \frac{\pi}{4} - 0 = \frac{\pi}{4}. \end{aligned}$$

3. Find  $\int_2^3 2xe^{-x^2} dx$

Ans.

$$\int_2^3 2xe^{-x^2} dx$$

{Let  $x^2 = u \Rightarrow 2xdx = du$ , when  $x = 2$ ,  $u = x^2 = 4$ ,

When  $x = 3$ ,  $u = x^2 = 9$ ,

So, lower limit changes to 4 and upper limit changes to 9}

$$= \int_4^9 e^u du$$

$$= [e^u]_4^9 = e^9 - e^4 \text{ (Ans)}$$

**Properties of Definite Integral**

---

1.  $\int_a^b f(x) dx = \int_a^b f(t) dt$

Explanation

Definite integral is independent of variable.

e.g.  $\int_2^3 x^2 dx = \int_2^3 u^2 du = \int_2^3 t^2 dt$

2.  $\int_a^b f(x) dx = - \int_b^a f(x) dx$

Explanation

If limits of definite integrals are interchanged then the value changes to its negative.

e.g.  $\int_2^3 x dx = - \int_3^2 x dx$

3.  $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$  where,  $a < c < b$ .

Explanation

If we integrate  $f(x)$  in  $[a, b]$  and  $c \in [a, b]$  such that  $a < c < b$ , then the above integral is same if we integrate  $f(x)$  in  $[a, c]$  and  $[c, b]$  and then add them.

e.g.  $\int_2^6 x dx = \int_2^4 x dx + \int_4^6 x dx$  verification

$$\int_2^6 x dx = \left[ \frac{x^2}{2} \right]_2^6$$

$$= \frac{6^2}{2} - \frac{2^2}{2} = \frac{36}{2} - \frac{4}{2} = 18 - 2 = 16 \text{ -----(1)}$$

$$\begin{aligned} \int_2^4 x dx + \int_4^6 x dx &= \left[ \frac{x^2}{2} \right]_2^4 + \left[ \frac{x^2}{2} \right]_4^6 = \left[ \frac{4^2}{2} - \frac{2^2}{2} \right] + \left[ \frac{6^2}{2} - \frac{4^2}{2} \right] \\ &= \left[ \frac{16}{2} - \frac{4}{2} \right] + \left[ \frac{36}{2} - \frac{16}{2} \right] = (8-2) + (18-8) \end{aligned}$$

$$= 6 + 10 = 16 \text{ ----- (2)}$$

From (1) and (2) we have,

$$\int_2^6 x dx = \int_2^4 x dx + \int_4^6 x dx \text{ (verified)}$$

4. 
$$\int_0^a f(x) dx = \int_0^a f(a-x) dx$$

e.g. 
$$\int_0^{\frac{\pi}{2}} \sin x dx = \int_0^{\frac{\pi}{2}} \sin(\frac{\pi}{2} - x) dx$$

verification

$$\begin{aligned} \int_0^{\frac{\pi}{2}} \sin x dx &= [-\cos \frac{\pi}{2} x]_0^{\frac{\pi}{2}} \\ &= -[\cos \frac{\pi}{2} - \cos 0] = -[0 - 1] = 1 \text{ ----- (1)} \end{aligned}$$

$$\int_0^{\frac{\pi}{2}} \sin(\frac{\pi}{2} - x) dx = \int_0^{\frac{\pi}{2}} \cos x dx$$

$$= [\sin x]_0^{\frac{\pi}{2}} = \sin \frac{\pi}{2} - \sin 0 = 1 - 0 = 1 \text{ ----- (2)}$$

From (1) and (2)

$$\int_0^{\frac{\pi}{2}} \sin x dx = \int_0^{\frac{\pi}{2}} \sin(\frac{\pi}{2} - x) dx \text{ (verified)}$$

5. (i) If f(x) is an even function, then

$$\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$$

(ii) If f(x) is an odd function, then

$$\int_{-a}^a f(x) dx = 0$$

Example: - By this formula without integration we can find the integral for

f(x) is an odd function if

$$f(-x) = -f(x)$$

Sinx, x, x<sup>3</sup> ..... are examples of odd functions. f(x) is

an even function if

$$f(-x) = f(x)$$

Cosx, x<sup>2</sup>, x<sup>4</sup> ..... are examples of even functions .

**Example:-**

$$\int_{-2}^2 x^2 dx = 2 \int_0^2 x^2 dx$$

{f(x) = x<sup>2</sup> is an even function as, f(-x) = (-x)<sup>2</sup> = x<sup>2</sup>. So, f(-x) = f(x)}

Similarly,

$$\int_{-a}^a f(x) dx = 0$$

Reason

$$f(x) = \sin x \Rightarrow f(-x) = \sin(-x) = -\sin x$$

$$\text{So, } f(-x) = -f(x)$$

□ f(x) is an odd function.

$$6. (i) \int_0^{2a} f(x) dx = 2 \int_0^a f(x) dx \quad \text{if } f(2a - x) = f(x)$$

$$(ii) \int_0^{2a} f(x) dx = 0 \quad \text{if } f(2a - x) = -f(x).$$

$$7. \int_a^b f(x) dx = \int_a^b f(a + b - x) dx$$

### Problems

**Q1.** Find  $\int_{-2}^1 |x| dx$

**Ans.**

$$|x| = \begin{cases} -x, & x < 0 \\ x, & x \geq 0 \end{cases}$$

|x| changes its definition at '0', so divide the integral into two parts (-2,0) and (0,1).

Now,  $\int_{-2}^1 |x| dx$

$$= \int_{-2}^0 |x| dx + \int_0^1 |x| dx \quad \{\text{Property (3)} \int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx\}$$

$$= \int_{-2}^0 -x dx + \int_0^1 x dx \quad \{\text{when, } -2 < x < 0 \text{ i.e. } x < 0 \text{ then, } |x| = -x \}$$

$$= - \left[ \frac{x^2}{2} \right]_{-2}^0 + \left[ \frac{x^2}{2} \right]_0^1 \quad \{\text{when, } 0 < x < 1 \text{ i.e. } 0 < x \text{ then, } |x| = x \}$$

$$= - \left[ \frac{0^2}{2} - \frac{(-2^2)}{2} \right] + \left[ \frac{1^2}{2} - \frac{0^2}{2} \right]$$

$$= - \left[ 0 - \frac{4}{2} \right] + \left[ \frac{1}{2} - 0 \right] = 2 + \frac{1}{2} = \frac{5}{2}$$

**Q2.**  $\int_{-6}^6 |x + 2| dx = ?$

**Ans.**

$$\int_{-6}^6 |x + 2| dx = \int_{-4}^8 |u| du$$

{Let u = x + 2 => du = dx, when, x = -6, u = -6 + 2 = -4}


$$\begin{aligned}
 & \{ \text{when, } x = 6, u = 6 + 2 = 8 \} \\
 & = \int_{-4}^0 |u| du + \int_0^8 |u| du \{ \text{property (3)} \} \\
 & = \int_{-4}^0 -u du + \int_0^8 u du \quad \{ \text{when } -4 < u < 0 \text{ then } |u| = -u \text{ and when } 0 < u < 8, \text{ then } |u| = u \} \\
 & = - \left[ \frac{u^2}{2} \right]_{-4}^0 + \left[ \frac{u^2}{2} \right]_0^8 \\
 & = - \frac{1}{2} [u^2]_{-4}^0 + \frac{1}{2} [u^2]_0^8 = - \frac{1}{2} [0^2 - (-4)^2] + \frac{1}{2} [8^2 - 0] \\
 & = - \frac{1}{2} (-16) + \frac{1}{2} (64) \\
 & = 8 + 32 = 40 (\text{Ans})
 \end{aligned}$$

### Area under plane curve

In our previous study we know that the definite integral represents the area under the curve.

### Area enclosed by curve and X- axis

Area enclosed by a curve  $y = f(x)$ , X-axis,  $x = a$  and  $x = b$  is given by



**Fig-2**

$$\text{Area} = \int_a^b f(x) dx$$

**Example – 1**

Find the area bounded  $y = e^x$ , X-axis  $x = 4$  and  $x = 2$

**Ans.**

Here  $y = e^x$  is the curve

Area of the curve bounded by X-axis,  $x = 4$  and  $x = 2$  is

$$\begin{aligned}\text{Area} &= \int_2^4 y \, dx = \int_2^4 e^x \, dx \\ &= [e^x]_2^4 = e^4 - e^2 \text{ (Ans)}\end{aligned}$$

**Example – 2**

Find the area enclosed by  $y = 9 - x^2$ ,  $y = 0$ ,  $x = 0$  and  $x = 2$ .

**Ans.**

$$\begin{aligned}\text{Area} &= \int_0^2 y \, dx = \int_0^2 (9 - x^2) \, dx \\ &= [9x - \frac{x^3}{3}]_0^2 = [ (9 \times 2) - \frac{2^3}{3} - (0-0) ] \\ &= 18 - \frac{8}{3} = \frac{54-8}{3} = \frac{46}{3} \text{ (Ans)}\end{aligned}$$

**UNIT-V**  
**DIFFERENTIAL EQUATION**

## Introduction

---

After the discovery of calculus, Newton and Leibnitz studied differential equation in connection with problem of Physics especially in theory of bending beams, oscillation of mechanical system etc. The study of differential equation is a wide field in pure and applied mathematics, physics and engineering.

## Objectives

---

1. Define differential equation.
2. Determine order and degree of differential equation.
3. Form differential equation from a given solution.
4. Solve differential equation by using different techniques.

## Definition of Differential Equation

---

A differential equation is an equation involving dependent variables, independent variables and derivatives of dependent variables with respect to one or more independent variables.

Here  $x$  is an independent variable,  $y$  is dependent variable and  $\frac{dy}{dx}$  is the derivative of the dependent variable w.r.t. the independent variable.

**Examples:** - i)  $\frac{dy}{dx} + xy = x^2$

ii)  $\frac{d^3y}{dx^3} + x^4 \frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^4 + y = \sin x$

iii)  $\frac{dy}{dx} + \sin x = \cos x$ .

Differential equations are of two types as follows: -

## Ordinary Differential Equation

---

An ordinary differential equation is an equation involving one dependent variables, one independent variable and derivatives of dependent variable with respect to independent variable. Mathematically  $F(x, y, \frac{dy}{dx}, \frac{d^2y}{dx^2}, \dots) = 0$

**Examples :** - i)  $\frac{dy}{dx} + y = x^2$

ii)  $\frac{d^3y}{dx^3} + x^4 \frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^4 + y = \sin x$

iii)  $\frac{dy}{dx} + y \tan x = \sec x$

## Partial Differential Equation

---

A partial differential equation is an equation involving dependent variables, independent variables and partial derivatives of dependent variable with respect to independent variables.

Examples:  $\frac{\partial z}{\partial x} + \frac{\partial z}{\partial y} = xy$

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = 0$$

In this chapter we only discuss about the Ordinary differential equation.

## Order and Degree of Differential equation

---

### Order

The order of the differential equation is the highest order of the derivatives occurring in it i.e. order of a differential equation is 'n' if the order of the highest order derivative term present in the equation is n.

Example 1:  $\frac{dy}{dx} + y = 2x$

The highest order derivative term in the equation is  $\frac{dy}{dx}$ , which has order 1.

∴ order of the differential equation is 1.

Example-2:  $-\frac{d^4 y}{dx^4} + \frac{d^3 y}{dx^3} + \left(\frac{dy}{dx}\right)^4 + y = x$

The highest order derivative term is  $\frac{d^4 y}{dx^4}$ , having order 4.

Hence the above differential equation has order 4.

### Degree

A differential equation is said to be of degree 'n', if the power i.e. highest exponent of the highest order derivative in the equation is 'n' after the equation has been freed from fractions and radicals as far as derivatives are concerned.

Before finding degree of a differential equation, first we have to eliminate those derivative terms present in fraction form i.e. in the denominator and derivatives with radicals i.e.

$$\sqrt{\frac{dy}{dx}}, \sqrt[3]{\frac{dy}{dx}}, \sqrt[4]{\frac{dy}{dx}} \text{ terms.}$$

**Example:-** Find the order and degree of following ordinary differential equations.

$$i) \frac{d^2y}{dx^2} = 3\left(\frac{dy}{dx}\right)^4 + x \quad ii) \left(\frac{d^4y}{dx^4}\right)^3 + \frac{d^3y}{dx^3} + 3\left(\frac{dy}{dx}\right)^4 + \cos x = 0$$

**Ans: - i)**  $\frac{d^2y}{dx^2} = 3\left(\frac{dy}{dx}\right)^4 + x$

Here  $\frac{d^2y}{dx^2}$  is the highest order derivative term.

Hence order of the differential equation is 2.

Again equation does not contain any derivative term in fractional form or with radical.

Power of the highest order derivative term  $\frac{d^2y}{dx^2}$  is 1.

Hence degree of differential equation is 1.

$$ii) \left(\frac{d^4y}{dx^4}\right)^3 + \frac{d^3y}{dx^3} + 3\left(\frac{dy}{dx}\right)^4 + \cos x = 0$$

From above it is clear that  $\frac{d^4y}{dx^4}$  is the highest order derivative term with power 3. Hence order =

4 and degree = 3

### Linear and Non-linear Differential Equation

---

A differential equation is said to be linear if it satisfies following conditions.

**i)** Every dependent variable and its derivatives have power '1'.

**ii)** The equation has neither terms having multiplication of dependent variable with its derivatives nor multiplication of two derivative terms.

Otherwise the equation is said to be non linear.

**Examples:-** i)  $\frac{dy}{dx} + xy = x^2$

ii)  $\frac{d^3y}{dx^3} + x^4 \frac{d^2y}{dx^2} + y = \sin x$

iii)  $\frac{d^3y}{dx^3} + y \frac{d^2y}{dx^2} = 4x$

iv)  $\left(\frac{dy}{dx}\right)^3 + 1 = 2 \frac{dy}{dx}$



$$v) \frac{d^3y}{dx^3} + \frac{dy}{dx} \frac{d^2y}{dx^2} + y = 4x^2$$

Among the above examples (i) and (ii) satisfy all the condition of linear equation. So the 1<sup>st</sup> two equations represent linear equations.

The (iii) is non linear because of the term  $y \frac{d^2y}{dx^2}$  which is a multiplication of dependent variable y and derivative term  $\frac{d^2y}{dx^2}$ .

The example (iv) is not linear due to the 1<sup>st</sup> term which contain  $\frac{dy}{dx}$  with power 3 violating the 1<sup>st</sup> condition of linearity.

The example (v) is not linear due to 2<sup>nd</sup> term which does satisfy the 2<sup>nd</sup> linearity property.

### Solution of a differential equation:-

---

The relationship between the variables of a differential equation satisfying the differential equation is called a Solution of the differential equation i.e  $y = f(x)$  or  $F(x,y)=0$  represent a Solution of the ordinary differential

equation  $F(x,y, \frac{dy}{dx}, \frac{d^2y}{dx^2}, \dots, \frac{d^ny}{dx^n}) = 0$  of order n if it satisfy it.

There are two types of Solutions i) General Solution ii) particular Solution

#### General Solution

---

The Solution of a differential equation containing as many arbitrary constants as the order of the differential equation is called as the general Solution.

Example- $y = A \cos x + B \sin x$  is a general Solution of differential equation  $\frac{d^2y}{dx^2} + y = 0$

#### Particular Solution

---

The Solution obtained by giving particular values to the arbitrary constants in the general solution is called particular solution

Example: -  $y = 3 \cos x + 2 \sin x$  is a particular Solution of differential equation  $\frac{d^2y}{dx^2} + y = 0$ .

### Differential equation of first order and first degree

---

A differential equation of 1<sup>st</sup> order and 1<sup>st</sup> degree involves x,y and  $\frac{dy}{dx}$ .

Mathematically it is written as  $\frac{dy}{dx} = f(x,y)$  or  $F(x, y, \frac{dy}{dx}) = 0$

## Solution of Differential equation of first order and first degree

---

The Solution of 1<sup>st</sup> order and 1<sup>st</sup> degree differential equation is obtained by following methods if they are in some standard forms as i) Variable separable form ii) Linear differential equation form.

### Variable Separable form

---

If the differential equation is expressed in the form,

$f(x)dy + g(y)dx = 0$ , then we say it variable separable form and this can be solved by integrating the terms separately as follows.

$$\text{Solution is given by } \int \frac{dy}{g(y)} = - \int \frac{dx}{f(x)}$$

$$\Rightarrow \log |g(y)| + \log |f(x)| = \log c$$

$$\Rightarrow g(y)f(x) = c$$

Where  $g(y)$  and  $f(x)$  are functions of  $y$  and  $x$  respectively, is called a variable and separable type equation.

**Example1:** - Solve  $\frac{dy}{dx} = x^2 + 2x + 5$

$$\text{Ans: } - \frac{dy}{dx} = x^2 + 2x + 5$$

$$\Rightarrow dy = (x^2 + 2x + 5)dx$$

Integrating both sides we have,

$$\Rightarrow \int dy = \int (x^2 + 2x + 5) dx$$

$$\Rightarrow y = \frac{x^3}{3} + \frac{2x^2}{2} + 5x + C = \frac{x^3}{3} + x^2 + 5x + c \text{ (Ans)}$$

**Example-2:** - Solve  $\frac{dy}{dx} = \frac{2y}{x^2+1}$ .

$$\text{Ans: } - \frac{dy}{dx} = \frac{2y}{x^2+1}$$

$$\Rightarrow \frac{dy}{2y} = \frac{dx}{x^2+1}$$

$$\Rightarrow \int \frac{dy}{2y} = \int \frac{dx}{x^2+1}$$

$$\Rightarrow \frac{1}{2} \log_e y = \tan^{-1}x + C$$

$$\Rightarrow \log_e y = 2 \tan^{-1}x + K \quad \{ 2C = K \text{ is a constant as } C \text{ is constant} \}$$

### Linear Differential Equation

---

A differential equation is said to be linear, if the dependent variable and its derivative occurring in the equation are of first degree only and are not multiplied together.

**Example:** -i)  $\frac{dy}{dx} + y = \sin x$  ii)

$$\frac{dy}{dx} + y \tan x = \sec x \text{ etc.}$$

### General form of linear differential equation

---

The general form of linear differential equation is given by,

$$\frac{dy}{dx} + Py = Q \text{ is linear in } y \text{ and } \frac{dy}{dx}.$$

Where P and Q are the functions of x only or constants.

This type of differential equation are solved when they are multiplied by a factor, which is called integrating factor (I.F.).

$$\mathbf{I.F.} = e^{\int P dx}$$

Then the solution is given by  $y (I.F.) = \int Q \cdot (I.F.) dx + C$ .

If equation is given in the form

$$\frac{dx}{dy} + Px = Q, \text{ where P and Q are functions of y only or constants and is linear in x and } \frac{dx}{dy},$$

then

$$\mathbf{I.F.} = e^{\int P dy}$$

Then the solution is given by  $x (I.F.) = \int Q \cdot (I.F.) dy + C$ .

This can be better understood by following examples.

**Example-11:** - Solve  $(1 + x^2) \frac{dy}{dx} + 2xy = x^3$  (2014-S, 2016-S, 2017-W).

**Ans:**  $-(1 + x^2) \frac{dy}{dx} + 2xy = x^3$

$$\Rightarrow \frac{dy}{dx} + \frac{2xy}{(1+x^2)} = \frac{x^3}{1+x^2}$$

By comparing with the general form of linear differential equation  $\frac{dy}{dx} + Py = Q$ .

Here  $P = \frac{2x}{(1+x^2)}$  &  $Q = \frac{x^3}{1+x^2}$

Now integrating factor I.F. =  $e^{\int P dx} = e^{\int \frac{2x}{(1+x^2)} dx}$  (putting  $1 + x^2 = t \Rightarrow 2x dx = dt$ )

$$= e^{\int \frac{dt}{t}} = e^{\ln t} = t = 1 + x^2$$

Solution is given by

$$y \times I.F. = \int Q.(I.F.)dx + C = \int \frac{x^3}{1+x^2}(1+x^2)dx + C = \int x^3 dx + c$$

$$\square y(1+x^2) = \frac{x^4}{4} + c$$

$$\therefore y = \frac{x^4}{4(1+x^2)} + \frac{c}{1+x^2}$$

**Example-12:** - Solve  $\frac{dy}{dx} + y = e^{-x}$

**Ans:** - By comparing the given equation with general form of linear differential equation we have,  $P = 1$  and  $Q = e^{-x}$

$$I.F. = e^{\int P dx} = e^{\int 1 dx} = e^x$$

Solution is  $y.(I.F.) = \int Q.(I.F.)dx + C$

$$= \int e^{-x}e^x dx + c$$

$$\Rightarrow ye^x = \int 1 dx + c = x + c$$

$$\Rightarrow y = xe^{-x} + ce^{-x}$$

**Example-13:** - Solve  $(1 - x^2)\frac{dy}{dx} - xy = 1$  (2017\_S)

**Ans:** -  $(1 - x^2)\frac{dy}{dx} - xy = 1$

$$\Rightarrow \frac{dy}{dx} - \frac{x}{1-x^2}y = \frac{1}{1-x^2}$$

Comparing with general form we have,

$$P = -\frac{x}{1-x^2} \text{ and } Q = \frac{1}{1-x^2}$$

Now I.F. =  $e^{\int P dx} = e^{\int -\frac{x}{1-x^2} dx}$  (Put  $1 - x^2 = t \Rightarrow -2x dx = dt \Rightarrow -x dx = \frac{dt}{2}$ )

$$= e^{\int \frac{dt}{2t}} = e^{\frac{1}{2} \ln t} = e^{\ln \sqrt{t}} = \sqrt{t}$$

$$= \sqrt{1 - x^2}$$

Solution is  $y.(I.F.) = \int Q.(I.F.)dx + C$

$$\Rightarrow y\sqrt{1-x^2} = \int \frac{1}{1-x^2}\sqrt{1-x^2} dx + C = \int \frac{dx}{\sqrt{1-x^2}} + c$$

$$= \sin^{-1}x + c$$

Hence solution of the differential equation is given by

$$y = \frac{\sin^{-1}x}{\sqrt{1-x^2}} + \frac{c}{\sqrt{1-x^2}} \quad (\text{Ans})$$

### Exercise

---

#### Question with short answers (2 marks)

1. Find the order and degree of the differential equations.

i)  $a \frac{d^2y}{dx^2} = [1 + (\frac{dy}{dx})^3]^{3/2}$  (2016-S)

ii)  $\frac{d^2y}{dx^2} = (\frac{dy}{dx})^{2/3}$  (2019-W)

2) Solve the following

i)  $\frac{dy}{dx} = e^{x+y}$  (2019-W)

ii)  $\frac{dy}{dx} = \frac{2y}{x^2+1}$  iii)  $\frac{dy}{dx} = \tan y$

#### Questions with long answers (5 marks)

3) Solve the following

i)  $(1 + x^2) \frac{dy}{dx} + 2xy - 4x^2 = 0$  (2019-W)

ii)  $\frac{dy}{dx} + y \tan x = \sec x$  (2015-S)